

# UTILITY PATENT APPLICATION TRANSMITTAL (Large Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.  
913.6600CIP

Total Pages in this Submission  
282

## TO THE ASSISTANT COMMISSIONER FOR PATENTS

Box Patent Application  
Washington, D.C. 20231

Transmitted herewith for filing under 35 U.S.C. 111(a) and 37 C.F.R. 1.53(b) is a new utility patent application for an invention entitled:

**COLD-SHOCK REGULATORY ELEMENTS, CONSTRUCTS THEREOF, AND METHODS OF USE**

and invented by:

Li Fang, Weining Jiang, Masanori Mitta, Masayori Inouye and Jean-Pierre Etchegaray

If a **CONTINUATION APPLICATION**, check appropriate box and supply the requisite information:

☒ **Continuation** ☐ **Divisional** ☐ **Continuation-in-part (CIP)** of prior application No.: PCT/US99/19030

Which is a:

☐ **Continuation** ☐ **Divisional** ☐ **Continuation-in-part (CIP)** of prior application No.: \_\_\_\_\_

Which is a:

☐ **Continuation** ☐ **Divisional** ☐ **Continuation-in-part (CIP)** of prior application No.: \_\_\_\_\_

Enclosed are:

### Application Elements

1. ☒ Filing fee as calculated and transmitted as described below
2. ☒ Specification having 77 pages and including the following:
  - a. ☒ Descriptive Title of the Invention
  - b. ☒ Cross References to Related Applications (if applicable)
  - c. ☐ Statement Regarding Federally-sponsored Research/Development (if applicable)
  - d. ☐ Reference to Microfiche Appendix (if applicable)
  - e. ☒ Background of the Invention
  - f. ☒ Brief Summary of the Invention
  - g. ☒ Brief Description of the Drawings (if drawings filed)
  - h. ☒ Detailed Description
  - i. ☒ Claim(s) as Classified Below
  - j. ☒ Abstract of the Disclosure

**UTILITY PATENT APPLICATION TRANSMITTAL**  
**(Large Entity)**

*(Only for new nonprovisional applications under 37 CFR 1.53(b))*

Docket No.  
913.6600CIP

Total Pages in this Submission  
282

**Application Elements (Continued)**

3. ☒ Drawing(s) *(when necessary as prescribed by 35 USC 113)*
- a. ☒ Formal Number of Sheets 29
- b. ☐ Informal Number of Sheets \_\_\_\_\_
4. ☒ Oath or Declaration
- a. ☐ Newly executed *(original or copy)* ☒ Unexecuted
- b. ☐ Copy from a prior application (37 CFR 1.63(d)) *(for continuation/divisional application only)*
- c. ☐ With Power of Attorney ☐ Without Power of Attorney
- d. ☐ DELETION OF INVENTOR(S)  
Signed statement attached deleting inventor(s) named in the prior application,  
see 37 C.F.R. 1.63(d)(2) and 1.33(b).
5. ☐ Incorporation By Reference *(usable if Box 4b is checked)*  
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied  
under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby  
incorporated by reference therein.
6. ☐ Computer Program in Microfiche *(Appendix)*
7. ☒ Nucleotide and/or Amino Acid Sequence Submission *(if applicable, all must be included)*
- a. ☒ Paper Copy
- b. ☒ Computer Readable Copy *(identical to computer copy)*
- c. ☒ Statement Verifying Identical Paper and Computer Readable Copy

**Accompanying Application Parts**

8. ☐ Assignment Papers *(cover sheet & document(s))*
9. ☐ 37 CFR 3.73(B) Statement *(when there is an assignee)*
10. ☐ English Translation Document *(if applicable)*
11. ☒ Information Disclosure Statement/PTO-1449 ☒ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Acknowledgment postcard
14. ☒ Certificate of Mailing
- ☐ First Class ☒ Express Mail *(Specify Label No.):* EL525816898US

# UTILITY PATENT APPLICATION TRANSMITTAL (Large Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.  
913.6600CIP

Total Pages in this Submission  
282

## Accompanying Application Parts (Continued)

15. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)

16. ☒ Additional Enclosures (please identify below):


International Search Report; Petition Under 37 CFR 1.84(b) with 16 photographs (in triplicate) and  
Petition Fee \$130.00; Submission of Formal Drawings;

## Fee Calculation and Transmittal

### CLAIMS AS FILED

For	#Filed	#Allowed	#Extra	Rate	Fee
Total Claims	56	- 20 =	36	x \$18.00	\$648.00
Indep. Claims	7	- 3 =	4	x \$78.00	\$312.00
Multiple Dependent Claims (check if applicable) <input type="checkbox"/>					\$0.00
BASIC FEE					\$690.00
OTHER FEE (specify purpose)					\$0.00
TOTAL FILING FEE					\$1,650.00

- ☒ A check in the amount of \$1,780.00 to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge and credit Deposit Account No. 13-3405 as described below. A duplicate copy of this sheet is enclosed.
- ☐ Charge the amount of \_\_\_\_\_ as filing fee.
- ☒ Credit any overpayment.
- ☒ Charge any additional filing fees required under 37 C.F.R. 1.16 and 1.17.
- ☐ Charge the issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

  
Signature

T. Daniel Christenbury, 31,750

Dated: 4/19/00

CC:

**Cold-Shock Regulatory Elements, Constructs Thereof, and Methods of Use**

This application is a continuation-in-part of Serial No. 09/293,427, filed April 16, 1999, entitled "Method and Constructs for Inhibiting Protein Expression in Bacteria," which is a continuation-in-part of Serial No. 08/769,945, filed December 19, 1996, entitled "Method and Constructs for Inhibiting Protein Expression in Bacteria," which is a continuation -in-part of Serial No. 08/203,806, filed March 1, 1994, entitled "Nucleic Acids Sequence, Stress-Induced Proteins and Uses Thereof," which corresponds to parent application Serial No. 07/852,013. This application also claims the priority of provisional application Serial No. 60/096,938, filed August 20, 1998, entitled "The 5'Untranslated Region of the Cold-Shock *cspA* Gene Regulates Translation Efficiency in Addition to mRNA Stability," and U.S. provisional application Serial No. 60/143,380, entitled "Translational Enhancement by an Element Downstream of the Initiation Codon in *Escherichia coli*."

**BACKGROUND OF THE INVENTION****1. Field of the Invention**

The invention relates to the control of bacterial gene expression, especially the regulation of bacterial gene expression under conditions of physiological stress. More specifically, the invention relates to the regulation of bacterial gene expression under conditions of physiological stress that induce the cold shock response of a bacterium.

**2. Description of the Related Art**

The regulation of bacterial gene expression occurs at many levels, including transcriptional control, or control of the synthesis of mRNA from a given gene; translational control, or the regulation of the efficiency by which the mRNA is translated into polypeptide sequence by the ribosome; and mRNA stability, or the efficiency at which a given mRNA

population within the cell is degraded and rendered inactive. The control of bacterial gene expression under conditions of physiological stress that elicit the cold shock response of the bacterium involve regulation at all three of the levels described above.

The response of bacteria to physiological stress involve the tightly controlled expression of a small number of genes that function to allow the cell to adapt to and function under stress conditions. For example, when bacterial cells are exposed to temperatures above the normal physiological temperature for that organism, a set of genes, designated the heat shock genes, are expressed. This response to elevated temperatures is well known and described in the prior art. Conversely, when bacterial cells are exposed to lower than physiological temperatures, a different set of genes, designated as cold shock (cs) genes, are expressed. Expression of the cs genes allow the cell to first adapt to the physiological stress, and subsequently grow under conditions of physiological stress. This invention relates to the specific processes that regulate the expression of cs genes.

When a culture of *Escherichia coli* is shifted from 37°C to 15 or 10°C, a number of proteins, called cold-shock proteins, are transiently induced during its growth lag period (Jones *et al.*, 1987; for review, see Thieringer *et al.*, 1998; Yamanaka *et al.*, 1998). *cspA*, consisting of 70 amino acid residues, has been identified as a major cold-shock protein (Goldstein *et al.*, 1990) and its three-dimensional structure has been determined by both X-ray crystallography (Schindelin *et al.*, 1994) and nuclear magnetic resonance spectroscopy (Newkirk *et al.*, 1994; Feng *et al.*, 1998) to consist of a five-antiparallel  $\beta$ -stranded structure. *cspA* can bind to single-stranded DNA and RNA without high sequence specificity and has been proposed to function as an RNA chaperone at low temperature (Jiang *et al.*, 1997).

To date, more than 50 proteins homologous to CspA have been identified in a large varieties of prokaryotes. Moreover, a region called cold-shock domain of eukaryotic Y-box

protein family, such as human YB-1 and *Xenopus* FRGY-2, shares more than 40% identity with *E. coli* *cspA* (for review, see Wolffe *et al.*, 1992), indicating that the cold-shock domain is well conserved throughout evolution. In *E. coli*, nine genes encoding *cspA*-like proteins, *cspA* to *cspI*, have been identified (for review, see Yamanaka *et al.*, 1998). Among them, 5 *cspA*, *cspB* and *cspG* are cold-shock inducible (Goldstein *et al.*, 1990; Lee *et al.*, 1994; Nakashima *et al.*, 1996) and interestingly, *cspD* is induced during stationary phase and upon nutrition starvation (Yamanaka and Inouye, 1997). It was proposed that the large *cspA* family of *E. coli* may have a function to respond to different environmental stresses (for review, see Yamanaka *et al.*, 1998).

10 *cspA* expression is transiently induced upon cold shock during the growth lag period called acclimation period. This period is considered to be required for cells to adapt to a new environmental condition. Indeed, during the acclimation period proteins involved in translation such as CsdA (Jones *et al.*, 1996), RbfA (Jones and Inouye, 1996) and CspA (Goldstein *et al.*, 1990) are specifically produced, which are considered to play important 15 roles in enhancing translation efficiency for non-cold-shock proteins at low temperature (for review, see Yamanaka *et al.*, 1998). Among these cold-shock proteins, *cspA* has been quite extensively investigated for the mechanism of its cold-shock induction (for review, see Yamanaka *et al.*, 1998). The *cspA* promoter is highly active at 37°C, even if CspA is hardly detected at this temperature (Fang *et al.*, 1997; Mitta *et al.*, 1997). Even if the *cspA* promoter 20 was replaced with the *lpp* promoter, a constitutive promoter for a major outer membrane protein, *cspA* expression is still cold-shock inducible (Fang *et al.*, 1997), indicating that the *cspA* induction at low temperature occurs mainly at levels of mRNA stability and its translation. The *cspA* promoter, however, contains an AT-rich upstream element (UP element) (Ross *et al.*, 1993) immediately upstream of the -35 region (Fang *et al.*, 1997;

Goldenberg *et al.*, 1997; Mitta *et al.*, 1997), which is considered to play an important role in efficient transcription initiation at low temperature. It has been demonstrated that the *cspA* mRNA becomes extremely stable upon cold shock, indicating that the mRNA stability plays a crucial role in cold-shock induction of *cspA* (Brandi *et al.*, 1996; Goldenberg *et al.*, 1996; Fang *et al.*, 1997).

An important and unique feature of the *cspA* mRNA is its unusually long 5'-untranslated region (5'-UTR) consisting of 159 bases (Tanabe *et al.*, 1992). This feature is also shared with other Class I cold-shock genes, which are dramatically induced after temperature downshift (for review, see Thieringer *et al.*, 1998), such as *cspB* (Etchegaray *et al.*, 1996) and *cspG* (Nakashima *et al.*, 1996). The 5'-UTR is considered to play a crucial role in the cold-shock induction of *cspA* (Brandi *et al.*, 1996; Jiang *et al.*, 1996; Goldenberg *et al.*, 1996; Bae *et al.*, 1997; Fang *et al.*, 1997; Goldenberg *et al.*, 1997; Mitta *et al.*, 1997).

Furthermore, it was recently shown that the 14-base downstream box (DB) located 12 bases downstream of the translation initiation codon of the *cspA* mRNA, which is partially complementary to a region called anti-downstream box of 16S rRNA (Sprengart *et al.*, 1996), plays an important role in efficient translation at low temperature (Mitta *et al.*, 1997). This region of the RNA sequence designated as the downstream box (DB) is complementary to bases 1469-1483 within the *E. coli* 16S rRNA (anti-DB sequence). It is speculated that formation of a duplex between the DB and anti-DB of 16S rRNA is responsible for translational enhancement (Sprengart *et al.*, 1996). The DB sequence has also been implicated in the translation of the  $\lambda$ cI mRNA, a mRNA that lacks any untranslated region and the SD sequence (Shean and Gottesman, 1992; Powers *et al.*, 1988). Interestingly  $\lambda$ cI translation was enhanced at 42°C in a temperature sensitive strain in which the amount of ribosomal protein S2 decreased at 42°C. It was proposed that the anti-DB sequence in S2

deficient ribosomes indirectly becomes more accessible to DB, resulting in enhancement of translation initiation of the  $\lambda$ CI mRNA. However, the role of the DB in  $\lambda$ CI translation initiation was disputed by Resch and coworkers (Resch et al., 1996). These authors constructed *lacZ* translational fusions with the  $\lambda$ CI gene to test the DB function. Since a  
5 deletion of 6 bases encompassing a portion of the DB sequence did not reduce the formation of the translation initiation complex, they disputed the existence of DB. Despite these elusive roles of DB (Sprengart and Porter, 1997) we have shown that the presence of a DB sequence in cold-shock mRNAs plays an important role in translation efficiency, and proposed that the DB is involved in the formation of a stable initiation complex at low temperature before the  
10 induction of cold-ribosomal factors (Mitta et al., 1997). Thus, *cspA* expression is regulated in a complex manner at levels of transcription, mRNA stability and translation.

Here, we constructed a series of deletion mutations in the 5'-UTR of *cspA* and analyzed their effects on *cspA* expression by examining the amount of mRNA, mRNA stability and translational efficiency. It was discovered that besides mRNA stability, the 5'-  
15 UTR plays a major role in translation efficiency of the *cspA* mRNA to enhance *cspA* expression upon cold shock. Specific regions of the 5'-UTR have been found to mediate the regulatory processes described above.



**BRIEF DESCRIPTION OF THE FIGURES**

Fig. 1 shows the plasmid pJG78 and the effects of the *cspA* upstream region on the chromosomal *cspA* expression and the synthesis of other cellular proteins: (A) pJG78. (B) the effects of the *cspA* upstream region on the chromosomal *cspA* expression and the synthesis of other cellular proteins.

Fig. 2 shows the prolonged expression of CspA and inhibition of cold-shock adaptation by pJG78 and pUC19-600.

Fig. 3 shows deletion analysis of the *cspA* upstream region for the *cspA* derepression function and inhibition of cold shock adaptation.

Fig. 4 shows the level of transcripts from the chromosomal and plasmid *cspA*.

Fig. 5 shows the requirement for the transcription of the 5' untranslated region of the *cspA* mRNA for the prolonged expression of *cspA* and inhibition of cold-shock adaptation.

Fig. 6 shows the effects of over-production of the 5' untranslated region of the *cspA* mRNA on the production of other cold-shock proteins and non-cold-shock protein.

Fig. 7 shows the effects of co-overproduction of *cspA* together with the 5' untranslated region of the *cspA* mRNA on the cold-shock response.

Fig. 8 shows the effects of overproduction of the first 25-base sequence of the *cspA* 5' UTR on CspA production.

Fig. 9. shows cold-shock induction of  $\beta$ -galactosidase: (A) Construction of *cspA-lacZ* fusions. The wild-type *cspA* is shown on the top. The *cspA-lacZ* fusion in each expression plasmid is shown from the 5' end of the *cspA* promoter upstream region to *lacZ*. Nucleotide numbers are given starting from the transcription initiation site as +1, determined by Tanabe *et al.*, 1992. The crossed hatched, open, dotted and slashed bars represent the *cspA* promoter, its 5' untranslated region, the *cspA* coding region and the *lacZ* coding region, respectively.

The solid boxes indicate the SD sequence. The positions of deleted regions are shown with nucleotide numbers. (B) Induction patterns of various deletion constructs. At mid-log phase, cultures of *E. coli* AR137 harboring various plasmids were shifted from 37°C to 15°C.

Samples were taken at 0, 1, 2, 3, 5, 7 and 10 h after the shift and  $\beta$ -galactosidase activity was measured. The *cspA-lacZ* fusions: pMM67 (○); pMM022 (●); pMM023 (□); pMM024 (■); pMM025 (Δ); and pMM026 (▲).

Fig. 10 shows the analysis of the mRNA stability. (A) Primer extension analysis of the cells harboring the *cspA-lacZ* fusions. At mid-log phase, cultures of *E. coli* AR137 harboring various plasmids were shifted from 37°C to 15°C. For measurement of the mRNA stability at 37°C, cultures were shifted back to 37°C after 30 min incubation at 15°C and rifampicin was added to the cultures to a final concentration of 200  $\mu$ g/ml. RNAs were extracted at 0 (lane 1), 1 (lane 2), 3 (lane 3) and 5 min (lane 4) after the addition of rifampicin. For measurement of the mRNA stability at 15°C, rifampicin was added 1 h after the temperature downshift, and then RNAs were extracted at 0 (lane 5), 5 (lane 6), 10 (lane 7), and 20 min (lane 8) after the addition of rifampicin. Primer extension was carried out. (B) Graphical presentation of the results shown in (A) for 37°C and 15°C, respectively. The radioactivities of transcripts were measured using a Phosphorimager and plotted using the transcript at zero time point as 100%: pMM022 (●); pMM023 (□); pMM024 (■); pMM025 (Δ); and pMM026 (▲).

Fig. 11 shows the analysis of the mRNA level and translational efficiency. (A) Primer extension analysis of the *cspA-lacZ* fusions. At mid-log phase, cultures of *E. coli* AR137 harboring various plasmids were shifted from 37°C to 15°C. RNAs were prepared from the culture at 37 °C (0 h; lane 1) and at 0.5 (lane 2), 1 (lane 3), 2 (lane 4), and 3 h (lane 5) after the temperature downshift. Primer extension was carried out as described previously (Mitta

*et al.*, 1997). (B) Graphical presentation of the relative amounts of mRNA. Relative mRNA amounts were calculated from the radioactivities of transcripts shown in (A) using the transcript of pMM67 at 37°C as 1. The relative amount is shown on the top of each column. Column 1, 0 h; column 2, after 0.5 h; column 3, after 1 h; column 4, after 2 h; and column 5, after 3 h. (C) Relative translational efficiencies of the *cspA-lacZ* mRNAs. Translational efficiencies at 15°C were calculated by dividing the increment of  $\beta$ -galactosidase activity during the first 2 h after cold shock by the amount of mRNA using the following formula:  $[(\text{Gal } 2 \text{ h}) \times (\text{OD } 2 \text{ h}) - (\text{Gal } 0 \text{ h}) \times (\text{OD } 0 \text{ h})] / (\text{avg. mRNA})$  where (Gal 0 h) and (Gal 2 h) are  $\beta$ -galactosidase activities at 0 h and 2 h after temperature downshift, respectively; (OD 0 h) and (OD 2 h) are the optical densities at 600 nm of the cultures at 0 h and 2 h after temperature downshift, respectively; (avg. mRNA) is the average of relative mRNA amounts at 0.5, 1 and 2 h after temperature downshift. Relative translational efficiency of each mRNA was calculated using the efficiency of mRNA of pMM67 as 100%. Column 1, pMM67; column 2, pMM022; column 3, pMM023; column 4, pMM024; column 5, pMM025; and column 6, pMM026.

Fig. 12 shows sequence similarities of *cspA*, *cspB*, *cspG*, and *cspI* mRNAs around the SD sequence and potential base pairing between *cspA* mRNA and 16S rRNA. Nucleotide numbers of *cspA* (Tanabe *et al.*, 1992), *cspB* (Etchegaray *et al.*, 1996), *cspG* (Nakashima *et al.*, 1996), and *cspI* mRNA (Wang *et al.*, 1999) are given starting from the major transcription initiation site as +1. The sequence of 16S rRNA is from Brosius *et al.*, 1978. Nucleotides identical in the three *csp* mRNAs are shown in bold letters. The 13-base homologous sequence in *cspA*, *cspB*, *cspG*, and *cspI* are boxed (the upstream box). Positions of the SD sequence and the initiation codon are underlined. Potential base pairings between *cspA*

mRNA and 16S rRNA are indicated by vertical lines. Positions of RNase V1 sensitive sites (Powers *et al.*, 1988) are dotted.

Fig. 13 shows the role of the 13-base upstream box sequence in the *cspA* 5'-UTR region in the *cspA-lacZ* expression. (A) Construction of *cspA-lacZ* fusions. The constructs are drawn in the same manner as shown in Fig. 9A except for a box with wavy lines, which represents the 13-base upstream box sequence, 5'-GCCGAAAGGCACA-3' (SEQ ID NO:48) located upstream of the SD sequence. pKNJ37 is identical to pMM007 except for the deletion of the 13-base sequence. In both pMM007 and pKNJ37, *cspA* was translationally fused to *lacZ*. pKNJ38 is identical to pKM67 (Mitta, *et al.*, 1997) except for the addition of the 13-base sequence by replacing the DNA fragment between *Xba*I and *Hind*III with synthesized oligonucleotides. In both pKM67 and pKNJ38, *cspA* was transcriptionally fused to *lacZ*. (B) Cold-shock induction of  $\beta$ -galactosidase. The *cspA-lacZ* fusions: pMM007 (○); pKNJ37 (●); pKM67 (Δ); and pKNJ38 (▲).

Fig. 14 shows a comparison of the secondary structures of the 5'-UTRs for the deletion constructs. Secondary structures of the 5'-UTR for each deletion construct was predicted with a nucleotide sequence analysis program (DNASIS-Mac; Hitachi Software Engineering Co. Ltd.) based on the method of Zuker and Stieger, 1982. Nucleotides are numbered as the position in the *cspA* mRNA starting from the transcription initiation site as +1. The position of the deletion in each mutant is shown by an arrow with the nucleotide numbers of the deleted region. The highly conserved 13-base sequence upstream of the SD sequence designated the upstream box are boxed. The initiation codon and the SD sequence are also boxed.

**Fig. 15** shows enhancement of *cspB* translation by DB. (A) *cspB*-DB-anti-DB complementarity: the *cspB*-DB sequence is boxed and encompasses the region from codons 5 to 9 (Mitta et al., 1997). Additional *cspB* mRNA-16S rRNA possible base pairings downstream of DB are also shown. The AUG codon is circled, the SD sequence is boxed and L-shaped arrows show the positions where the *cspB* gene was fused to *lacZ*. (B) Translational *cspB-lacZ* fusion constructs. On the top, the *E. coli cspB* gene is depicted from its 5' end. In pB3, pB13 and pB17, the *lacZ* gene is fused to *cspB* at residue +177 (3 aa), +200 (13 aa) and +212 (17 aa), respectively. The pB13sd and pB17sd are the same as pB13 and pB17, respectively, except that their SD sequences are changed from 5'-AGGA-3' to 5'-CTTC-3'.

(C)  $\beta$ -galactosidase activity of the *cspB-lacZ* constructs obtained before (time 0) and after (1, 2 and 3 hr) temperature shift from 37 to 15°C. *E. coli* AR137 cells were transformed with pB3, pB13, pB13sd, pB17 and pB17sd were grown in medium, and at mid-log phase ( $OD_{600} = 0.4$ ) cultures were shifted from 37 to 15°C.  $\beta$ -galactosidase activity was measured. (D) mRNA levels of pB3, pB13, pB17 or pB13sd after temperature shift from 37 to 15°C: the *cspB-lacZ* mRNAs were detected by primer extension before temperature downshift (time 0) and at 1, 2 and 3 hrs after temperature shift. (E) mRNA stability from pB3, pB13, pB17 and pB13sd: *E. coli* AR137 cells transformed with pB3, pB13, pB17 and pB13sd were grown under the same conditions described above. At mid log phase, the culture was shifted to 15°C and after 30 min., rifampicin was added to a final concentration of 0.2 mg/ml (time 0). Total RNA was extracted at 5, 10 and 20 min. after rifampicin addition. The *cspB-lacZ* mRNAs were detected by primer extension.

**Fig. 16** shows the effect of a perfectly matching DB enhancing the translation of *cspA*. (A) Translational *cspA-lacZ* fusion constructs. The *cspA* gene structure from its 5'-end is showed at the top. pJG78DB1 and pJG78DB2 were constructed from pJG78 as described

in Experimental Procedures. The DB sequences of pJIG78DB1 (12 matches) and pJIG78DB2 (15 matches) are shown at the bottom.

(B)  $\beta$ -Galactosidase activity of the *cspA-lacZ* fusion constructs after cold shock at 15°C. *E. coli* AR137 cells transformed with pJIG78, pJIG78DB1 or pJIG78DB2 were grown in LB medium, and at mid-log phase ( $OD_{600}=0.4$ ) cultures were shifted from 37°C to 15°C.  $\beta$ -galactosidase activity was measured before (time 0) and 1, 2 and 3 hr after the shift.

(C) Detection of the *cspA-lacZ* mRNAs. Total RNA from *E. coli* AR137 cells carrying pJIG78, pJIG78DB1 or pJIG78DB2 was extracted at the same time points indicated above and used as a template for primer extension. (D) mRNA stability from the *cspA-lacZ*

constructs. *E. coli* AR137 cells transformed with pJIG78, pJIG78DB1 and pJIG78DB2 were grown as described above. At mid-log phase, the cultures were shifted to 15°C and after 30 minutes rifampicin was added to a final concentration of 0.2 mg/ml (time 0). Total RNA was extracted at 5, 10 and 40 minutes after rifampicin addition. The *cspA-lacZ* mRNAs were detected by primer extension.

Fig. 17 shows that a perfectly matching DB enhances translation at 37°C: (A) pIN-*lacZ* constructs. The *XbaI-SaII* fragment from pJIG78 or pJIG78DB2 was inserted into the *XbaI-SaII* sites of pIN-III to create pINZ and pINZDB1, respectively which then were used to create pINZDB2, pINZDB3 and pINZDB4.

(B)  $\beta$ -Galactosidase activity of the pINZ-*lacZ* constructs. Cultures of *E. coli* AR137 cells transformed with pINZ, pINZDB1, pINZDB2, pINZDB3 and pINZDB4 were grown at 37°C under the same conditions described in Figure 1. IPTG (1 mM) was added at mid-log phase

to each culture.  $\beta$ -Galactosidase activity was measured before (time 0) and at 0.5, 1, 2 and 3 hr after IPTG addition.

(C) mRNA sequences of the pINZ-*lacZ* constructs showing the position of SD, AUG and DB.

The *lacZ* in pJJG78 has a 10-match DB. The perfect match DB located after the 5th codon

has 16 residues complementary with the anti-DB. (D) shows the rate of  $\beta$ -galactosidase

synthesis of the pINZ-*lacZ* constructs. Cultures of *E. coli* AR137 cells carrying pINZ or

pINZDB1 were grown at 37°C under the same conditions described above. IPTG (1 mM)

was added at mid-log phase to each culture. Rate of  $\beta$ -galactosidase synthesis was measured

before (time 0) and 0.5, 1, 2, 3 and 4 hr after IPTG addition. Cells were pulse-labeled with

trans-[<sup>35</sup>S]-methionine. Cell extracts from each time point were analyzed by 5% SDS-PAGE

and the  $\beta$ -galactosidase synthesis was measured by phosphorimager. The ratio of  $\beta$ -

galactosidase synthesis of pINZ and pINZDB1 is shown at each time point.

Fig. 18 shows ribosomal fractionation of *E. coli* JM83 cells transformed with pINZ or

pINZDB1. Ribosomal particles were isolated as described by Dammel and Noller, 1995.

Cultures of *E. coli* JM83 cells carrying pINZ or pINZDB1 were grown at 37°C in LB medium

containing 50 mg/ml of ampicillin. At mid-log phase ( $OD_{600}=0.4$ ) 1 mM of IPTG was added

to each culture. Chloramphenicol (0.1 mg/ml) was added at 15, 30 and 60 min after IPTG

addition. The cell extracts were then layered on top of a 5-40% (w/w) sucrose gradient. The

polysomes and ribosomal subunits were separated by centrifugation at 151,000 x g for 2.5 hr

at 4°C. The polysome profiles were then detected by using a FPLC system. 0.2 ml from each

fraction (0.5 ml) were spotted on a Nitrocellulose membrane using the Minifold II Slot-Blot

System (Schleicher and Schuell). The *lacZ* mRNA was detected by hybridization using the

[<sup>32</sup>P]-labeled M13-47. Phosphorimager values from the hybridization are plotted at the right

side. The pINZ and pINZDB1 mRNAs are shown in closed and open squares, respectively.

**Fig. 19.** shows translational enhancement by a perfectly matching DB at 37°C. **(A)**

Estimation of pINZ and pINZDB1 mRNAs. Cultures of *E. coli* JM83 carrying pINZ or pINZDB1 were grown at 37°C under the same conditions described in Figure 4. Total RNA extracted at 15, 30 and 60 min after IPTG (1 mM) addition was used as a template for primer

5 extensions according to the procedure described previously. **(B)**  $\beta$ -Galactosidase activity of pINZ and pINZDB1 in multi-copy expression system. *E. coli* JM83 cells transformed with pINZ or pINZDB1 were grown at 37°C under the same condition described in Figure 20A.  $\beta$ -Galactosidase activity was measured before (time 0) and 0.5, 1, 1.5, 2 and 2.5 hr after IPTG (1 mM) addition (open circles and squares). Closed circles and squares represent the activities in the absence of IPTG.

**Fig. 20** shows cell-free synthesis of  $\beta$ -galactosidase from pINZ and pINZDB1. **(A)** pINZ or pINZDB1 DNA (160 ng; 1  $\mu$ l) was added to the *E. coli* 30S extract (20 $\mu$ l) (Promega) and the transcription-translation coupled reaction was carried out. Lane 1, pINZ DNA; lane 2, pINZDB1 DNA; and lane 3, a control reaction without added DNA. Samples were  
15 precipitated with acetone and analyzed by 15% SDS-PAGE to detect the production of  $\beta$ -galactosidase. **(B)** Time course *in vitro* synthesis of  $\beta$ -galactosidase from pINZ and pINZDB1 was carried out as described above. Samples were taken after 15, 30, 60 and 120 min incubation at 37°C. **(C)** Each reaction from the time course experiment described above was done in duplicate with non radioactive methionine, spotted on nitrocellulose membrane  
20 and hybridized with [<sup>32</sup>P] labeled M13-47 oligonucleotide.

**Fig. 22** shows translational enhancement of pINZDB1 in cells with S2-depleted ribosomes **(A)**  $\beta$ -galactosidase activity from pINZ and pINZDB1. *E. coli* CS240 and CS239 (Shean and Gottesman, 1992) were transformed with pINZ or pINZDB1 and cultures were grown at 30°C in LB medium. At mid-log phase the cells were shifted to 42°C in the presence of 1 mM of



5 IPTG.  $\beta$ -Galactosidase activity was measured as Miller units before (time 0) and at 0.5, 1, 1.5, 2.5 and 3.5 hr after shift to 42°C. (B) Relative induction of the *lacZ* expression between pINZDB1 and pINZ in cells with S2-depleted ribosomes. Before the shift to 42°C (time 0) the ratio of the  $\beta$ -galactosidase expression from pINZ and pINZDB1 in CS239 to that of CS240 was estimated as 1, and the ratios after the shift to 42°C were calculated accordingly.

### SUMMARY OF THE INVENTION

10 The invention comprises an isolated nucleic acid molecule that prolongs the expression of cold-shock inducible genes under conditions of physiological stress that elicit the cold-shock response in bacteria.

The invention further comprises an isolated nucleic acid molecule that represses the expression of cold-shock inducible genes under physiological conditions.

15 The invention further comprises an isolated nucleic acid molecule that enhances translation of cold-shock inducible genes under conditions of physiological stress that elicits a cold shock response in bacteria.

20 What has been discovered is that three of the essential processes that govern the regulation of genes that are expressed under conditions of physiological stress that elicit the cold shock response of a bacterium are mediated by the 5'-untranslated region (5'-UTR) of the mRNA transcripts encoding cold shock inducible proteins. It has been discovered that specific regions of the 5'-UTR of cold shock inducible mRNA transcripts mediate (A) the transient nature of cs gene expression after exposure of the cells the physiological stress that induces the cold shock response, (B) the repression of cs gene expression at physiological temperature, and (C) enhancement of the translation of cold shock inducible mRNA transcripts by the ribosome.

It has also been discovered that DNA constructs encoding these regulatory elements, either alone or in combination with themselves, or with other regulatory sequences known in the art, e.g., promoters, downstream boxes and transcriptional terminator sequences, can be exploited to prolong the expression and increase the efficiency of translation of cs genes or of a heterologous gene under conditions of physiological stress that induce the cold shock response of a bacterium, and repress the expression of said genes under normal physiological condition. This is advantageous for the hyper-expression of a desired protein product which may be unstable at physiological temperatures, or may fold improperly and exist in inclusion bodies within the host cell at physiological temperatures, or may be degraded by the host cell.

The invention offers a further advantageous trait in that the translation of normal cellular proteins is inhibited under the conditions where the desired protein product is being hyper-expressed. This allows the desired product to accumulate in the host cell and simplifies the purification of the desired product from host cell polypeptides.

Accordingly, the invention also comprises transformed bacteria carrying the vectors which contain one or more of the above described elements and transformed bacteria carrying vectors with one or more of the above described elements and a target gene sequence for expression. Such target sequence can be a cold-shock gene sequence or a heterologous gene sequence.

#### **DETAILED DESCRIPTION OF THE INVENTION**

As reported by Sprengart et al., the downstream box (DB) of bacteria plays an important role in the translation of mRNA to produce proteins. The DB binds to a portion of the bacterial 16S rRNA near the 3' end and is thought to help position the mRNA and rRNA in proper relative position for translation to occur.

It has been discovered that during the time when the ADB is annealed to the DB of an overexpressed mRNA, the 16S rRNA is not capable of participation in the translation of cellular mRNAs other than the annealed overexpressed mRNA. It has been further discovered that the entire protein-making machinery of a bacterium may be shut down by providing to the bacterium an mRNA, which encodes a DB which is substantially complementary to the ADB of the 16S rRNA, which anneals to all or substantially all of the bacterial 16S rRNA.

The term "complementary" is used herein, it is intended to include "substantially complementary". Thus, the term "complementarity" does not require perfect complementarity. It is sufficient that the two sequences be "complementary" as defined in Kahl, Dictionary of Gene Technology, VCH Publishers, Inc. (1995), which is incorporated herein by reference. That is, two nucleotide sequences are complementary if they are capable of forming a hydrogen-bonded duplex with each other according to Watson-Crick base-pairing rules. Two complementary RNA sequences, or an RNA and a DNA sequence, will form pairings of A-U, G-C, or G-U. "Complete complementarity" is not required.

"Homologous" as used herein refers to molecules which have substantially the same molecular sequence of the referenced nucleic acid sequence, but may contain additions, deletions, or substitutions. Homologous molecules are defined as those molecules which hybridize under low or high stringency conditions to a nucleic acid molecule that is precisely complementary to the referenced nucleic acid molecule and which performs the same function as the referenced nucleic acid.

By way of example, and not limitation, low stringency conditions for hybridization are: Filters containing DNA are pretreated for 6 hours at 40°C in a solution of 35% formamide, 5xSSC, 50 mM Tris-HCL (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1%

BSA, and 500 µg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and  $5\text{--}20 \times 10^6$  cpm  $^{32}\text{P}$ -labeled probe. Filters are then incubated in hybridization mixture for 18-20 hours at 40°C, and washed for 1.5 hours at 55°C in a solution containing 2xSSC, 25 mM Tris-HCL (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash solution is replaced with fresh wash solution and incubated for an additional 1.5 hours at 60°C. The filters are blotted dry and exposed for autoradiography. If desired, or if required, a third wash step may be conducted for 1.5 hours at a temperature of 65-68°C and the filters can be reexposed to film.

For example, and not by way of limitation, high stringency conditions may be as follows: Prehybridization of filters containing the nucleic acid to be probed is carried out for 8 hours to overnight at 65°C in buffer containing 6xSSC, 50 mM Tris-HCL (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA and 500 µg/ml denatured salmon sperm DNA. The filters are hybridized for 48 hours at 65°C in a prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and  $5\text{--}20 \times 10^6$   $^{32}\text{P}$ -labeled probe. Washing of filters is performed for 1 hour in a solution of 2xSSC, 0.01 PVP, 0.01% Ficoll, and 0.01% BSA at 37°C, followed by a wash for 45 minutes in 0.1xSSC at 50°C. The filters are blotted dry and exposed for autoradiography. Other protocols for high and low stringency hybridization known in the art may be substituted.

As is known, the ADB is a nucleotide sequence of about 14 bases which is positioned in the 3' end of the 16S rRNA, in close proximity to the decoding region of 16S rRNA. The 16S rRNA nucleotide sequence of known bacteria is known and can be found in the GenBank database. Thus, for a selected bacterium, the ADB can be readily identified by comparison to the sequence of the ADB in a bacterium in which the sequence is known, for example *E. coli*. Once the ADB is identified, a DB complementary to the ADB can be constructed, and incorporated into an appropriate mRNA, as described below.

The mRNA of the invention is an isolated mRNA or an mRNA which has been transcribed from an isolated DNA. The mRNA comprises an initiation codon, which codon is preferably AUG. Other suitable initiation codons for the mRNA include GUG and UUG.

The mRNA of the invention further comprises a downstream box sequence, which is typically 3' to the initiation codon. The codons of the DB may or may not be in-frame with the initiation codon. The DB sequence may be immediately adjacent to the initiation codon so that there are no intervening nucleotides. Generally, the DB is separated from the initiation codon by an intervening nucleotide sequence between 1 and 30 nucleotides long. The base sequence of the intervening sequence is immaterial and may be constituted of any sequence of nucleotides. Preferably, the intervening nucleotide sequence is 9 to 15 nucleotides in length, with a most preferred length of 12 nucleotides. Alternatively, the DB may overlap the initiation codon. That is, any one of the three nucleotides of the initiation codon of the mRNA of the invention may form the 5' end of the DB.

The DB sequence of the mRNA of the invention is a nucleotide sequence which is complementary to the ADB of the 16S rRNA of a bacterium. Generally, the DB is between 6 and 20 bases long, preferably between 8 and 14 bases long, although the DB may be longer than 20 bases. For example, the DB may comprise nucleotides which are complementary to nucleotides 3' or 5', or both, to the ADB. Regardless of length of the DB, a higher degree of complementarity between the DB and the ADB is associated with more effective annealing, resulting in more efficient inhibition of bacterial protein synthesis, in accordance with the method of the invention.

In addition to the initiation codon, the DB, and any intervening sequence, the mRNA construct of the invention may comprise a nucleotide sequence 5' to the initiation codon or 3' to the DB. For example, the mRNA construct may comprise a sequence 3' to the DB which

encodes a polypeptide or may comprise a termination codon. Likewise, the mRNA construct may comprise an untranslated sequence and/or a Shine-Dalgarno sequence 5' to the initiation codon.

The length of the mRNA construct, including the initiation codon, any intervening sequence, and DB, and exclusive of any additional nucleotides at the 5' or 3' end, may be any length between 8 nucleotides to about 45 nucleotides. Of course, if the mRNA comprises a 5' or 3' sequence in addition to the above essential components, such as a Shine-Dalgarno sequence, the mRNA may be much longer, up to several hundreds of nucleotides in length.

Preferably, although not necessarily, the mRNA construct is free of sites for RNA endonucleases. It is especially preferred that the portion of the mRNA construct comprising the essential portions of the construct, that is the initiation codon and the DB, be free of sites for RNA endonucleases, which might otherwise degrade the mRNA construct and free the bacterial 16S rRNA to bind to bacterial mRNAs.

The mRNA construct of the invention may have a sequence which is similar or identical to an mRNA sequence found naturally in a bacterium. For example, the mRNAs for several cold-shock proteins, such as the mRNAs for *E. coli* proteins CspA, CspB, CspG, CsdA, and RbfA, comprise a Shine-Dalgarno sequence, an initiation codon, and a downstream box substantially complementary to the ADB of the *E. coli* 16S rRNA. Other *E. coli* mRNAs which contain a Shine-Dalgarno sequence, an initiation codon, and a downstream box complementary to the *E. coli* ADB include RecA, Hns, NusA, InfB, and CspD.

Below are several non-limiting examples of suitable DBs for the mRNA construct. Each of the following DB is substantially complementary to the ADB of the *E. coli* 16S rRNA which ADB has the sequence:

ADB 3' (-1481) UACUUAGUGUUUCA (-1469) 5' (SEQ ID NO:1)

DB #1: 5' AUGACUGGUAUCGU 3' (SEQ ID NO:2)

DB #2: 5' AUGACUGGUUUCGU 3' (SEQ ID NO:3)

DB #3: 5' AUGACUGGUUUAGU 3' (SEQ ID NO:4)

5 DB #4: 5' AUGAGUUAUGUAGA 3' (SEQ ID NO:5)

DB #5: 5' AUGGCAGAAAAGAAU 3' (SEQ ID NO:6)

A suitable mRNA construct according to the invention can be constructed using any one of the above DBs, or other suitable DB, for example:

5' AUGX<sub>(n)</sub>AUGACUGGUAUCGU 3' (SEQ ID NO:7)

10 where n is a whole number from 0 to 30, and X is G, C, U, or A, wherein each occurrence of X may be the same as or different from any other occurrence of X. Alternatively, the 5' end of the DB overlaps the initiation codon.

The DNA of the invention is any isolated DNA which encodes for a mRNA which is suitable for the mRNA construct of the invention, as described above. The DNA may further  
15 comprise an additional nucleotide sequence 5' to the initiation codon, which sequence may include a promoter sequence. Such promoter sequences may be used to control transcription of the mRNA construct. The DNA may comprise a sequence 5' to the initiation codon which sequence has a function other than as a promoter, such as a Shine-Dalgarno sequence, and/or a sequence which has no known function. The DNA may comprise a sequence 3' to the  
20 portion encoding the DB of the mRNA construct, which sequence may include, for example, a termination codon, or may encode a polypeptide, and a sequence required for transcription termination.

An example of a suitable DNA which encodes for the mRNA construct of the invention is: 5' ATGY<sub>(n)</sub>ATGACTGGTATCGT 3' (SEQ ID NO:8) where n is a whole number

from 0 to 30, and Y is G, C, T, or A, wherein each occurrence of Y may be the same as or different from any other occurrence of Y. Alternatively, the 5' end of the DB overlaps the initiation codon, ATG. The DNA may contain additional sequences, as stated above, at the 5' and/or 3' end of the DNA.

5           The DNA sequence of the invention may be contained within a vehicle or cloning vector, such as in a plasmid or phage vector. The DNA sequence in the vector may be under the control of a promoter sequence located 5' to the initiation codon. These vectors containing the DNA of the invention may be used to transform a host bacterium which may be used to overexpress the mRNA of the invention, that is to produce the mRNA in the  
10       bacterium at levels higher than produced in similar non-transformed bacteria. Any bacterium which may be transformed by means of a cloning vector is a suitable host for the DNA sequence of the invention. Methods of producing cloning vectors and transforming bacteria are known in the art and are taught, for example, in Ausubel et al., Current Protocols in Molecular Biology, J. Wiley & Sons, Inc. (1995), which is incorporated herein by reference.

15           Overexpression of the mRNA sequence of the invention results in the production of the mRNA in an amount which is higher than that found normally in the bacteria. To whatever extent the mRNA is overexpressed, the production of bacterial proteins is inhibited. If the mRNA is expressed at a high enough level, production of bacterial proteins will be completely stopped, which may lead ultimately to death of the bacterium.

20           Therefore, the construct producing the mRNA is useful as an antibiotic to kill or to stop the growth of bacteria. The construct producing the mRNA may be packaged in a bacteriophage which would permit the mRNA to be used as a disinfectant or as a topical antibiotic preparation. It is conceivable that strategies for delivery will be devised to permit transformation of bacteria which are causing infection of a plant or animal, such as a mammal



like humans, dogs, cats, cattle, horses, and livestock. Such antibiotics are safe for use in eukaryotes, as eukaryotes lack the 16S rRNA that is present in bacteria.

According to the method of the invention, a mRNA comprising an initiation codon and a DB which is complementary to the ADB of the 16S rRNA of a bacterium, is caused to be overexpressed in a bacterium, and is then allowed to anneal to the ADB of the 16S rRNA of the bacterium, thereby inhibiting production of proteins encoded by other mRNAs in the bacterium.

Any means of delivery which results in overexpression of the mRNA of the invention is suitable for the method of the invention. For example, the bacterium may be transformed by means of a vehicle harboring a DNA sequence which codes for the mRNA of the invention.

If desired, expression of the mRNA sequence of the invention is controlled by placing the DNA sequence under the control of an inducible promoter. For example, if it is desired to kill a harmful bacterium or block its growth while sparing a beneficial bacterium, the DNA sequence may be placed under the control of a promoter which is responsive to a product which is present only in the first bacterium. In this way, the lethal antibiotic effect of the mRNA of the invention will affect only the undesirable, harmful bacterium.

Another means of controlling the expression of the protein production-inhibiting mRNA sequence is to employ a DNA sequence which codes for an mRNA which is unstable under certain conditions.

For example, the 5' untranslated region (5' UTR) of the mRNA of the *E. coli* cold-shock protein, CspA, contains a region immediately 5' to the Shine-Dalgarno region which is susceptible to degradation, presumably by RNase E, at physiological growth temperatures of about 37°C. Therefore, the *cspA* mRNA containing the 5' UTR is unstable under normal

growth conditions, having a half life estimated to be approximately 12 seconds. Other cold-shock proteins, such as *E. coli* CspB and CsdA, are similarly unstable at physiological growth temperatures due to instability of their mRNAs. Upon cold shock, such as when the temperature is reduced to 15°C, the half life of the *cspA* mRNA increases dramatically, to about 15 minutes, an increase in stability of about 75 times over the mRNA at normal physiological growth temperatures.

Because of the instability at 37°C of an mRNA containing the 5' UTR of *cspA* mRNA, this region, or the 5' UTR of the *cspB* or *csdA* mRNA, can be used to control the expression of the mRNA sequence of the invention, so that its antibiotic effect occurs only below physiological growth temperatures, such as under cold-shock conditions. The antibiotic effect of the method of the invention is augmented at cold-shock conditions because a cold-shocked bacterium requires new ribosomal factors, whose synthesis is blocked by overproduction of an mRNA containing the DB sequence.

The antibiotic effect of the method of the invention in which the mRNA of the invention is caused to be overexpressed within a bacterium is increased concomitantly with an increase in copy number of the mRNA which is to be expressed. That is, whereas a minimal overexpression of the mRNA of the invention will inhibit the production of proteins by the bacterium, such an inhibition may not be sufficient to prevent further growth of the bacterium or to kill the bacterium. Higher levels of expression of the mRNA result are positively correlated with increased inhibition of protein production. When the copy number is sufficiently high in the bacterium, protein production will be completely blocked.

A similar effect is noted with respect to complementarity of the DB of the overexpressed mRNA and the ADB of the bacterial 16S rRNA. Overexpression of an mRNA comprising a DB with 100% complementarity will be more efficient in binding to the ADB

than will be an mRNA comprising a DB with lesser (75%) complementarity. Thus, the protein blocking effect of an mRNA having a more highly complementary DB will be more pronounced compared to that of an mRNA having a less complementary DB. Therefore, when using an mRNA having a less complementary DB, it may be useful to express the mRNA in a higher copy number to achieve the same or similar antibiotic results as with an mRNA having a more complementary DB.

The translational inhibitory properties of the downstream box are also advantageous for overexpressing a heterologous gene in a transformed bacterium after cold shock. Inhibition of the translation of endogenous bacterial proteins will allow the heterologous gene product to accumulate to very high levels in the transformed organism. Furthermore, a construct containing the downstream box in conjunction with a strong promoter and the 5' untranslated region of a cold shock inducible gene, which functions to stabilize the mRNA transcript at reduced temperature, will direct efficient high level expression of the heterologous gene at reduced temperature.

Another important embodiment of the invention relates to the role of the 5' -end untranslated region of the mRNA for *cspA*, the major cold-shock protein of *E. coli*, in cold shock adaption.

When the culture temperature of exponentially growing *E. coli* cells are shifted from 37 to 10°C, there is a growth lag period before reinitiation of cell growth (Jones et al. 1987). Similar to the heat-shock response, *E. coli* responds to the temperature downshift by inducing a specific pattern of gene expression called cold-shock response, which includes induction of a set of proteins defined as cold-shock proteins (Jones et al. 1992; for review, see Jones and Inouye 1994). The cold-shock response occurs during the lag period of cell growth, and is considered to be required for cellular adaptation to low temperature.

CspA, the major cold-shock protein in *E. coli*, is dramatically induced upon temperature downshift, whose production reaches as high as 13% of total protein synthesis (Goldstein et al. 1190). Interestingly, however, CspA production during cold-shock response is transient and drops to a basal level at the time of reinitiation of cell growth at low temperature. CspA consists of 70 amino acid residues, and shows 43% identity to the “cold-shock domain” of the eukaryotic Y-box protein family which is known to be associated with gene regulation and mRNA masking (for review, see Wolffe et al. 1992; Wolffe 1993). The three-dimensional structure of CspA has been determined, consisting of five anti-parallel  $\beta$ -sheets which form a  $\beta$ -barrel structure (Newkirk et al. 1994; Schindelin et al. 1994). Two RNA binding motifs, RNP1 and RNP2, are identified on  $\beta$ 2 and  $\beta$ 3 sheets, respectively. In the structure, seven out of eight aromatic residues are located on the same surface and a single-stranded DNA was shown to interact with these surface aromatic residues (Newkirk et al. 1994). It has been proposed that CspA function as an RNA chaperone to facilitate translation efficiency at low temperature (for review, see Jones and Inouye 1994).

### Cold Box

In one embodiment of the invention, a DNA sequence is capable of prolonging the normally transient expression of the cold shock genes during the adaptation of a bacterium to physiological stress that elicits the cold shock response. Thus, the transient nature of the cold shock response and the normally transient expression of the cold shock inducible gene is blocked and expression is continued at high levels for an extended period of time, such as at least 2 to 3 hours. The DNA sequence that is competent to confer this activity comprises the 5'-UTR or at least a portion of the 5'UTR of a cold shock inducible mRNA transcript and a promoter, active under conditions of physiological stress that induce the cold shock response

in a bacterium. Furthermore, it was found that not the entire 5'-UTR was essential, but that the sequence responsible for blocking the transient expression of the cold shock genes, like *cspA*, resides within the first twenty five nucleotides of the 5'-UTR.

Comparison of this nucleotide sequence which is competent for this activity with other cold shock genes showed that *cspB*, *cspG* and *csdA* possessed similar sequences within their respective 5'-UTR and were expressed in a transient manner in response to physiological stress that induces the cold shock response. This suggests that these genes are regulated in a similar manner to *cspA*. Based upon these sequence comparisons, the sequence responsible for this activity, hereafter designated the cold box, was shown to be situated between nucleotides +1 and +11 of the 5'-UTR of *cspA*, *cspB*, *cspG* and *csdA*.

The cold box normally functions to down regulate the expression of cold shock genes. This was shown by hyper-expression of just the *cspA* 5'-UTR in *E. coli*, which resulted in a prolongation of the expression of cold shock inducible genes in these cells. The cold box functions probably by interacting with the CspA protein itself, or with another protein whose function is dependent upon CspA. This was shown by subsequently hyper-expression of the CspA protein in the *E. coli* cells described above and observing a suppression of the prolonged expression. Therefore, the cold box appears to comprise a repressor binding site that functions to repress the expression of cold shock genes after their induction, resulting in transient expression of these genes. As the levels of CspA protein increase after induction of its gene, the CspA begins to interact with the cold box, either directly by binding the cold box sequence or indirectly by stimulating another factor to interact with the cold box, resulting in repression of the cold shock inducible genes.

**Repressor of cold shock gene expression at physiological temperatures**

The 5' UTR is unique is that it is longer than most *E. coli* 5'UTRs. It plays multiple functions in *cspA* expression. It represses *cspA* expression at 37°C. These different activities have been mapped by Yamanaka et al (unpublished manuscript) by constructing and analyzing defined deletions through the 5'-UTR region. The sequences mediating repression of *cspA* expression at 37°C reside between +56 and +117 of the 5'-UTR of *cspA* (see Yamanaka *et al.*, unpublished manuscript). It is likely that optimal repression requires other sequences in the 5'-UTR as well. In other words, the effect of deleting the +56 to +117 region was significant, but did not account for the entire level of repression observed with the intact 5'-UTR. Thus, his conclusion that other sequences may also play a role in repressing *cspA* expression at 37°C. However, the localization of at least a portion of the repression activity to a specific region of the 5'-UTR is novel. The 5'-UTR is starting to be dissected into functional domains and the +56 to +117 region represents a new functional region.

The UTR also has a positive effect on mRNA stability at 15°C. This was determined by noting an increase in steady state levels of mRNA of constructs having the UTR. However, even though more mRNA was present in these experiments, the mRNA was not translated unless the DB was present.

The effects of increasing mRNA stability and increasing mRNA translatability are thus mediated by independent sequences of the *cspA* transcript. However, the specific sequences that mediate mRNA stability are not well characterized in this region.

## Translational enhancement of cold shock mRNA

The 5'-UTR enhances the translatability of *cspA* transcripts through a sequence situated between +117 and +143 of the 5' UTR. Comparison of this region between different cold shock inducible revealed a 13 base sequence (+123 - +135) having the sequence 5'-

GCCGAAAGGCACA-3' (SEQ ID NO:48) that was conserved among *cspA*, *cspB*, and *cspG* and may represent enhancer of translation. Thus, this sequence mediates efficient translation of the mRNA that possesses it. This 13 base region exhibits homology with the 16S rRNA, similar to what was previously observed with the DB, but the new translational enhancer region is complementary to a different region of the 16S rRNA than DB. Therefore, this region of the 5'-UTR may also assist in translating *csp* mRNAs by a mechanism similar to the downstream box interactions with ribosomal RNA.

In another embodiment of the invention, expression plasmids, capable of high level expression of cold shock inducible genes, or of a heterologous gene, are constructed. Such expression plasmids contain DNA fragments encoding the 5'UTR or a portion or portions thereof of a cold shock inducible gene, comprising one or more of the regulatory elements described above, positioned downstream of a promoter that is functional under conditions of physiological stress that induce the cold shock response in a bacterium. Such promoters may be selected from the group including the *cspA*, *cspB*, *cspG* or *csdA* promoters, the *E. coli lpp* promoter, or any other such promoter that is active under conditions of physiological stress. Such expression plasmids may also contain downstream of the promoter and DNA fragment encoding the 5'-UTR comprising one or more of the regulatory elements described, and a transcriptional terminator sequence. Examples of transcriptional terminator sequences are well known in the art and include sequences such as the *E. coli rrnB* terminator.

The expression plasmids of the invention may also comprise a restriction site, or a sequence comprising multiple restriction sites, such site or sites situated between the 5'-UTR encoding one or more of the regulatory elements and the transcriptional terminator to facilitate the insertion of a heterologous gene.

Finally, the expression plasmids of the invention may comprise additional sequences

known in the art to facilitate the efficient translation of the expressed gene. Such sequences may include a Shine-Dalgarno sequence, situated between the 5'UTR sequence and the restriction site(s) and/or a DNA fragment encoding a downstream box, situated between the Shine-Dalgarno sequence and the restriction site(s). The source of the Shine-Dalgarno sequence is not especially limited, and may be derived from cold shock proteins or may be from another gene. Such expression plasmids are capable of directing high level expression of a heterologous gene for a prolonged period of time under conditions of physiological stress that elicit the cold shock response of a bacterium. Under these conditions, the synthesis of endogenous proteins by the host bacterium is blocked, allowing the product of the heterologous gene to accumulate to high levels within the cell.

In a preferred embodiment of the invention, the expression vector comprises a promoter, a 5'UTR, cold box, Shine-Dalgarno sequence and a downstream box (DB). This expression vector may be used so that a nucleic acid molecule encoding a target protein may be ligated into the expression vector using at least one restriction site.

In another embodiment of the invention, nucleic acid molecules encoding proteins that are unstable, or that fold improperly in the bacterial host cell at physiological temperatures can be expressed at temperatures below the physiological temperature of the host bacterium. Such nucleic acid molecules may be inserted, preferably in-frame to the restriction site or sites to allow transcription of the nucleic acid and translation of the resulting mRNA. Such conditions may facilitate the proper folding, increase the stability or decrease the rate of degradation of the expressed product.

The plasmids or vectors may be used to transform bacteria by any method known in the art, including, but not limited to calcium chloride transformation, electroporation, and the like. The bacterial species which may be transformed is not particularly limited. In a



preferred embodiment, *E. coli* is used. The transformed bacteria may be used to overexpress a target protein of interest, or may be used to produce large amounts of the plasmids or vectors for subsequent isolation and purification.

In a preferred embodiment for overexpression of a target protein, a nucleic acid molecule encoding a target protein is ligated into the plasmid using one or more restriction sites, preferably in-frame to the initiation codon, if an initiation codon is provided upstream of the insertion site for the target protein. Alternatively, if the nucleic acid molecule encoding the target protein has its own initiation codon, the nucleic acid encoding the target protein may be ligated into the plasmid such that its own initiation codon will serve as the initiation codon in the transcript.

The constructs may be designed and assembled to include a selectable marker. That is, for example, a drug resistance gene which allows transformed bacteria to grow in the presence of a drug which does not permit the growth of non-transformed bacteria. Any selectable marker known in the art may be used. Examples of selectable markers include, but are not limited to ampicillin resistance, neomycin resistance, kanamycin resistance and tetracycline resistance.

The constructs may also include an inducible promoter. The inducible promoter may direct the synthesis of the target protein upon induction. For example, and not by way of limitation, a lacZ promoter may be included which may be induced by the addition of IPTG to the culture medium.

In a preferred embodiment, the constructs will contain a promoter which is active under conditions that elicit a cold-shock response in bacteria, such as by shifting the temperature of the culture medium containing the transformed bacteria to 10-15°C. Preferably, overexpression of a target protein in conditions that elicit a cold-shock response

reduces the synthesis of at least one native bacterial protein. More preferably, the synthesis of many native proteins is reduced or blocked.

The following examples are provided to illustrate aspects of the invention and are not to be construed in any way as limiting the scope of the invention, which is defined in the appended claims.

## EXAMPLES

### Example 1: Construction of Plasmids

pJIG02 was constructed from pJIG01 (Goldstein *et al.*, 1990) as follows: A 998-bp fragment which contains the entire *cspA* gene was obtained from pJIG01 by *Hind*III and *Xmn*I digestion. This fragment was then treated with the Klenow fragment of DNA polymerase (Life Technologies), and inserted into the *Sma*I site of pUC9.

pJIG21 was constructed from pJIG02 by creating an *Xba*I site immediately upstream of the Shine-Dalgarno sequence of *cspA* as follows:

+13AATTI(A)C(T)TAG(A)AGGTAA+153 (SEQ ID NO:9)(the original nucleotides in the parentheses were substituted by the underlined nucleotides; ref. 1). pJIG81 was constructed from pJIG02 by creating an *Xba*I site immediately downstream of the transcription initiation site of *cspA* as follows: +1ACGGTTCTAGACGTA+15 (SEQ ID NO:10)(nucleotides underlined represent the inserted bases).

pJIG78 is a transcriptional fusion of the 0.6-kb *cspA* upstream region and *lacZ* as follows: the 1-kb *Eco*RI/*Bam*HI fragment containing *cspA* from pJIG21 was filled in with Klenow enzyme and ligated into the *Sma*I site of pUC19. Then, the 0.6-kb *Xba*I fragment containing the *cspA* regulatory region (from -457 to +143) was excised and ligated into the *Xba*I site in pKM005 (Inouye, M. *et al.*, 1983) in the correct orientation.

pUC19-600 was constructed by insertion of the 0.6-kb *EcoRI/XbaI* fragment from pJIG21 into the *EcoRI/XbaI* sites of pUC19. pJIG81/X,S containing fragment 1 (Figure 3) was constructed by removing the 0.74-kb *XbaI/SalI* fragment from pJIG81. Both ends were treated with Klenow fragment, followed by self-ligation. All the other constructs shown in Fig. 3 were made by PCR (Boehringer Mannheim protocol). PCR amplified fragments were inserted into the *SmaI* site of pUC19. All PCR products were confirmed by DNA sequencing (Sanger *et al.*, 1977).

p2JTEK was constructed as follows: PCR product by primer 3549 5'-CGGCATTAAGTAAGCAGTTG-3' (SEQ ID NO:11) and primer 4428 5'-CTGGATCCTTAATGGTCTGTACGTCAAACCGT-3' (SEQ ID NO:12) was cloned into the *SmaI* site of pUC19. This PCR product contains *ospA* from -146 to +25 as the *ospA* transcription start site is defined as +1. Then the transcriptional terminator of *ospA* was amplified by PCR using primer 6290 5'-CGGAATTCAGCCTGTAATCTCT-3' (SEQ ID NO:13) and 4860 5'-CTGTCGACTTACTTACGGCGTTGC-3' (SEQ ID NO:14). The PCR product was then digested with *EcoRI* then cloned into the plasmid described above which was digested with *EcoRI* and *SspI*. The 52-bp *KpnI* and *EcoRI* fragment from Bluescript II SK was then cloned into the *EcoRI* and *KpnI* site. All PCR products were confirmed by DNA sequencing (Sanger *et al.*, 1977).

p6mTEK was constructed in the same way as p2JTEK except that the first PCR was carried out with different primers: primer 3552 5'-GACAGGATTAATAAATCGAG-3' (SEQ ID NO:15) and 6196 5'-AACCGTTGATGTGCA-3' (SEQ ID NO:16). This PCR product encompasses *ospA* from -278 to +6 as the *ospA* transcription start site is defined as +1. All PCR products were confirmed by DNA sequencing (Sanger *et al.*, 1977).

The pulse-labeling experiments were carried out as described previously (Jiang *et al.*,

1993). Proteins were analyzed either by polyacrylamide SDS-gel electrophoresis (Inouye S. *et al.*, 1982) or by two-dimensional electrophoresis as described previously (Jones *et al.*, 1987).

Each 5'-UTR deletion was introduced by two step PCR. For the first step, two PCRs were carried out for each mutation. One reaction was done with a combination of primer

5 67F, 5'-ccttgtagCCGATTAATCATAAATATG-3' (SEQ ID NO:17)(nucleotides -67 to -49 of *cspA*), and mutation primer R, which contains desired mutated sequence and is complimentary to *cspA*, and the other was done with primer #4311,

5'-ccggatccagGTTGAACCATTTT-3' (SEQ ID NO:18) (complementary to +186 to +198) and mutation primer F, which also contains the same mutation and is same direction to *cspA*. The

10 lower cases are extra nucleotides to create *NheI* or *BamHI* site (underlined) and the numbers are given using major transcription initiation site at +1, which was determined by Tanabe *et al.* (1992). For construction of pMM022, pMM023, pMM024, pMM025, and pMM026,

primer D1R, 5'-ACTACACT/TTGATGTGCATTAGC-3' (SEQ ID NO:19)(complementary to -15 to +1 / +28 to +35), primer D2R, 5'-CAACGATAA/GCTTTAATGGTCTGT-3' (SEQ ID

15 NO:20) (complementary to +13 to +27 / +56 to +64), primer D3R,

5'-TAAAGG/CTCTGAAGGGACTT-3' (SEQ ID NO:21)(complementary to +41 to +55 / +86 to +91), primer D4R, 5'-CGGCGATAT/AATGTGCACTACGAGGG-3' (SEQ ID NO:22) (complementary to +69 to +85 / +118 to +126), and primer D5R,

5'-TACCTTTAA/GGCGTGCTTTACAGATT-3' (SEQ ID NO:23)(complementary to +101 to

20 +117 / +144 to +152) was used as the mutation primer R and primer D1F,

5'-GCACATCA/AGTGTAAGGCAA-3' (SEQ ID NO:24)(nucleotide -8 to +1 / +28 to +42), primer D2F, 5'-TAAAGC/TTATCGTTGATACCC-3' (SEQ ID NO:25)(nucleotide +22 to +27 / +56 to +70), primer D3F, 5'-TCAAGAG/CCTTTAACGCTTCAAAA-3' (SEQ ID NO:26)(nucleotide +49 to +55 / +86 to +102), primer D4F,

5'-GCACATT/ATATCGCCGAAAGGC-3' (SEQ ID NO:27)(nucleotide +79 to +85 / +118 to +132), and primer D5F, 5'-AAAGCAGCC/TTAAAGGTAATACACT-3' (SEQ ID NO:28)(nucleotide +108 to +117 / +144 to +159), was used as the mutation primer F, respectively, where the position of each deletion is indicated by a slash. A plasmid, pJG02 (Goldstein et al. (1990), which contains the wild-type *cspA*, was used as template DNA. Then, each set of the first PCR products were mixed, heat denatured, annealed, and extended by Taq polymerase. The resulting products were further amplified by PCR using primer 67F and primer #4311. The final PCR products were digested with *NheI* and *BamHI*, and inserted into the *XbaI*-*BamHI* site of pKM005 (Inouye 1983). For pKNJ37, the PCR fragment was cloned into the *XbaI*-*BamHI* site of pRS414X, a pRS414 derivative (Simons *et al.*, 1987), in which the unique *SmaI* site has been changed to an *XbaI* site.

For the construction of pMM007, PCR was carried out using primer 67F and primer #4311 as primers and pJG02 as a template. The PCR fragment was digested with *NheI* and *BamHI*, and inserted into the *XbaI*-*BamHI* site of pRS414X.

pKNJ38 was constructed as follows: oligonucleotide #8509,

5'-

CTAGCCGAAAGGCACAAATTAAGAGGGTATTAATAATGAAAGGGGGAATTCCA-3' (SEQ ID NO:29), and oligonucleotide #8510,

5'-AGCTTGGAATCCCCCTTCATTATTAATACCCTCTTAATTTGTGCC'TTCGG-3' (SEQ ID NO:30) were first annealed and then cloned into pKM67 (Mitta *et al.*, 1997) digested with *XbaI* and *HindIII*.

The DNA sequences of all the constructs were confirmed by DNA sequencing using the chain-termination method (Sanger *et al.*, 1977).

*E. coli* AR137 harboring different plasmids was grown at 37°C to mid-log phase in 15

ml of M9-Casamino acid medium using a 125-ml flask. The culture was then transferred to a 15°C shaking water bath. Culture temperature reached 15°C from 37°C within 2 to 3 min under the condition used. A 1.5-ml culture was taken immediately before the temperature downshift (0hr), and at 1, 2, 3, 5, 7, and 10 hr after the temperature downshift.

- 5     β-galactosidase activity of the culture was measured according to Miller (1972). The assay was done in duplicate at each time points.

*cspB* DNA fragments were amplified by PCR using synthetic oligonucleotide primers containing the *Bam*HI site at the 5' end. A plasmid, pSJ7 (Lee *et al.*, 1994) carrying the wild-type *cspB* gene was used as a template DNA to create the PCR fragments B3, B13 and B17.

- 10    The 5'-end oligonucleotide primer used in each of the above PCR reactions is 5'-CCGGATCCAGCTTTAATATAGCT-3' (SEQ ID NO:31). The 3'-end oligonucleotide primers for the PCR products B3, B13, and B17 were 5'-CCGGATCCAGATTGACATTCTACA-3' (SEQ ID NO:32), 5'-CCGGATCCAGGTAAACCATTTT-3' (SEQ ID NO:33), and
- 15    5'CCGGATCCAGACCTTTATCAGCGTT-3' (SEQ ID NO:34), respectively. A deletion of the SD sequence in the *cspB* gene was created by site-directed mutagenesis using the QuickChange™ Site Directed Mutagenesis Kit (Stratagene). The PCR reaction was carried out using the pSJ7 plasmid as a template and the oligonucleotides Bsd1 (5'-GAAAGGCTCAAGTTACTTCATGTAGAATG-3')(SEQ ID NO:35) and Bsd2 (5'-
- 20    CATTCTACATGAAGTAACTTGAGCCTTTC-3')(SEQ ID NO:36) to create pSJ7sd. pSJ7sd was used as a template to make the PCR fragments B13sd and B17sd. The 5' and 3' end oligonucleotide primers used in these PCR reactions are the same as the ones used for the PCR fragments B13 and B17. All of the above PCR products were cloned at the *Bam*HI site of pRS414 vector (Simmons *et al.*, 1987; Lee *et al.*, 1994) to create the pB3, pB13, pB13sd,

pB17, and pB17sd constructs.

The *cspA-lacZ* fusion constructs were made by the insertion of annealed oligonucleotides at the *EcoRI* site of the pJG78 (Jiang *et al.*, 1996). Annealed oligonucleotides DB1 (5'-AATTAATCACAAAGTGGG-3') (SEQ ID NO:37) with DB1' (5'-AATTCCCACCTTTGTGATT-3') (SEQ ID NO:38) or DB2 (5'-AATTATGAATCACAAAGTGGG-3') (SEQ ID NO:39) with DB2' (5'-AATTCCCACCTTTGTGATTCAT-3') (SEQ ID NO:40) were used to create pJG78DB1 or pJG78DB2 constructs, respectively.

The pIN-*lacZ* constructs were made by inserting the *XbaI-SalI* fragments from pJG78 or pJG78DB2 into the *XbaI-SalI* sites of pIN-III (Jiang *et al.*, 1996) to create pINZ and pINZDB1, respectively. Then, the annealed oligonucleotides ZDB2 (5'-CTAGCCCTT-ATTAATAATGAAAGGGGAATTATGAATCACAAAGTGGG-3') (SEQ ID NO:41) with ZDB2' (5'-AATTCCCACCTTTGTGATTCATAATTCCCCCTTCATTATTAATAAGGG-3') (SEQ ID NO:42) were inserted at the *XbaI-EcoRI* sites of pINZ to create pINZDB2. Annealed oligonucleotides ZDB3 (5'-CTAGCCCTTATTAATAATGAATCACAAAGTGGG-3') (SEQ ID NO:43) with ZDB3' (5'-AATTCCCACCTTTGTGATTCATTATTAATAAGGG-3') (SEQ ID NO:44) or ZDB4 (5'-CTAGAGGG-TATTAATAATGAATCACAAAGTGGG-3') (SEQ ID NO:45) with ZDB4' (5'-AATTCCCACCTTTGTG-ATTCATTATTAATACCCT-3') (SEQ ID NO:46) were inserted at the *XbaI-EcoRI* sites of pINZ to construct pINZDB3 and pINZDB4, respectively.

#### $\beta$ -Galactosidase Activity

*E. coli* AR137 (*pcnB*<sup>-</sup>) (Harlocker *et al.*, 1993) or JM83 (*pcnB*<sup>-</sup>) harboring different plasmids were grown at 37°C to mid-log phase in 20 ml of LB medium containing 50  $\mu$ g/ml

of ampicillin in a 125 ml flask. The cultures were then transformed to a 15°C shaking water bath or isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) was added to a final concentration of 1 mM. A 100 ml culture was taken at each time point.  $\beta$ -galactosidase activity was measured according to Miller's procedure (Miller, 1972).

5

### Example 2: Isolation of RNA, primer extension and ribosome analysis

*E. coli* AR137 harboring different plasmids was grown under the same conditions used for the  $\beta$ -galactosidase assay described above. In order to estimate the amount of *cspA-lacZ* mRNA, a 1.5-ml culture was taken at each time point and RNA was extracted by the hot-phenol method as described by Sarmientos et al., 1983.

10

For the mRNA stability experiments at 15°C, rifampicin was added at 1 hr after the temperature downshift at a final concentration of 200  $\mu$ g/ml to stop transcription, and a 1.5-ml culture was taken at each time point. For the mRNA stability at 37°C, the culture was first shifted to 15°C for 30 min to accumulate the mRNAs. Then, a 5-ml culture was taken and mixed with a 5 ml of the medium containing 400  $\mu$ g/ml rifampicin in a glass flask kept in 37°C shaking water bath. The medium was prewarmed at 60°C. Using this method, the culture temperature immediately changed to 32°C and reached 37°C within 1 min.

15

The *cspA-lacZ* mRNAs were detected by the primer extension method as described previously (Jiang et al. 1993) using a  $^{32}$ P-labeled primer M13-47, 5'-CGCCAGGGTTTCCAGTCACGAC-3' (SEQ ID NO:47), which is complementary to a 5'-end coding sequence of *lacZ*. The products were separated on a denatured polyacrylamide gel (6%) and quantified by the use of Phosphorimager (BioRad).

20

### Pulse Labeling



Cultures of *E. coli* AR137 (*pcnB*<sup>-</sup>) cells carrying pINZ or pINZDB1 were grown at 37°C under the same conditions used for the  $\beta$ -galactosidase assay. IPTG (1 mM) was added at mid-log phase to each culture. At each time point, 1 ml of the culture was labeled for 5 min with 100  $\mu$ Ci of trans-[<sup>35</sup>S]methionine (1,175 Ci/mmol) (NEN Life Science Products) as described previously (Jones *et al.*, 1996). Cell extracts from each time point were loaded on a 5% SDS-PAGE and  $\beta$ -galactosidase synthesis was measured by phosphorimager.

### Ribosome Isolation

Cultures of *E. coli* JM83 (*pcnB*<sup>+</sup>) cells carrying pINZ or pINZDB1 were grown in 600 ml of LB medium in a 4-liter flask under the same condition as described above. At mid-log phase IPTG (1 mM) was added to each culture. Ribosomal particles were isolated by the procedure described by Dammel and Noller (1995), with some modifications: An 100-ml aliquot from the original culture was taken at each time point and chloramphenicol was added to a final concentration of 0.1 mg/ml to stop cell growth. Cells were immediately collected by centrifugation (5,000 x g for 10 min at 4°C), resuspended in buffer I [20 mM Tris-HCl (pH 7.6), 10 mM MgCl<sub>2</sub>, 100 mM NH<sub>4</sub>Cl, 6 mM  $\beta$ -mercaptoethanol and 1 mg/ml lysozyme] and frozen at -80°C for a few hours. The cells were lysed by the freeze-thaw method (Ron *et al.* 1996). The cell extracts (0.5 ml) were then layered on top of a 5-40 % (w/w) sucrose gradient (7.5 ml) and the polysomes and ribosomal subunits were separated by centrifugation at 151,000 x g for 2.5 hr at 4°C using a Beckman SW-41 rotor. The polysome profiles were detected by an FPLC system and a total of 15 fractions of 0.5 ml each were collected.

### Detection of the *lacZ-mRNA*

From each polysome fraction (0.5 ml) 0.2 ml was spotted on a Nitrocellulose membrane using the Minifold® II Slot-Blot System (Schleicher and Schuell). The *lacZ*-mRNA was detected by hybridization using the [<sup>32</sup>P]-labeled M13-47 primer (S. Inouye and M. Inouye, 1991) and the amount of *lacZ*-mRNA was estimated by phosphorimager.

5

### ***In vitro* Translation**

Using the *E. coli* S30 Extract System for Linear Templates Kit (Promega), the transcription-translation coupled reaction was carried out according to the manufacturer's protocol as follows; To 20 µl of Pre-mix containing all the amino acids except for methionine, 10 µCi of trans-[<sup>35</sup>S]methionine (1,175 Ci/mmol; NEN Life Science Products) and the *E. coli* S30 extracts, 160 ng of pINZ or pINZDB1 (1 µl) was added and the mixture was incubated at 37°C for 45 min. The products were precipitated with acetone and analyzed by 15 % SDS-PAGE.

### **EXAMPLE 3: Multicopy effects of the *cspA* upstream region on cold-shock adaptation.**

It has been shown that the *cspA* gene is induced immediately after the temperature downshift from 37°C to either 15 or 10°C and that the rate of CspA production reaches a peak after 1 hr at 15°C and 2 hr at 10°C after the temperature shift (Goldstein *et al.*, 1990). After this time point, CspA production sharply drops to a new basal level. The period of this transient production of CspA corresponds to the duration of growth arrest, known as the lag period, which is observed after cold shock (Jones *et al.*, 1987). Thus, such a transient expression of *cspA* is considered to be required for cellular adaptation to lower temperatures.

In order to characterize this transient expression of *cspA*, we attempted to identify the region required for the regulation of the *cspA* expression during the adaptation period. For

this purpose, pJG78 was first constructed, in which the 600-bp *cspA* upstream region was transcriptionally fused to the *lacZ* gene (Fig. 1A). This 600-bp upstream region of *cspA* encompasses the region from -457 to +143 which is right before the Shine-Dalgarno sequence of *cspA*, as the *cspA* transcription initiation site is defined +1 (Goldstein *et al.*, 1990). *E. coli* strain CL83 was transformed with pJG78 and the production of  $\beta$ -galactosidase was examined by pulse-labeling cells with [<sup>35</sup>S]-methionine at 0, 0.5, and 3 hr after temperature downshift from 37 to 15°C. As controls, CL83 cells alone as well as CL83 cells transformed with vector pKM005 (Inouye 1983) were also used. As shown in Fig. 1B, for both CL83 and CL83/pKM005 the expression of *cspA* was highly induced at 0.5 hr after the temperature downshift (Fig. 1B, lanes 2 and 5, respectively). However, as shown previously (Goldstein *et al.*, 1990), this high expression is transient and reduced to a new basal level at 3 hr (Fig. 1B, lanes 3 and 6, respectively). Note that no *cspA* expression was detected at 0 time point (Fig. 1B, lanes 1 and 4, respectively) and that  $\beta$ -galactosidase was not produced at any time point for both strains (Fig. 1B, lanes 1 to 6).

In contrast to CL83 and CL83/pKM005,  $\beta$ -galactosidase was clearly induced in the cells with pJG78 upon the temperature downshift (Fig. 1B, lanes 7 to 9), indicating that the 600-bp upstream region of *cspA* is sufficient for the cold-shock induction. Surprisingly, the production of *cspA* was no longer transient but remained at a high level even 3 hr after cold shock in the cells harboring pJG78 (Fig. 1B, compare lane 9 with lanes 3 and 6). Since pJG78 does not contain the *cspA* coding sequence, the high production of CspA at 3 hr after temperature downshift is attributed to the chromosomal *cspA* gene. It appears that under the conditions used, the chromosomal *cspA* gene failed to be repressed, in other words it became derepressed. Interestingly, there is another band indicated by X in Fig. 1B, whose expression pattern was almost identical to that of *cspA*. It is a cold-shock protein and its production was

also derepressed in the presence of pJG78. This cold-shock protein X has been recently identified as CsdA which associates with ribosomes (Jones *et al.*, 1994).

It should also be noted that the synthesis of most cellular proteins was blocked to a larger extent in the cells harboring pJG78 at low temperature than that in the CL83 cells and  
5 CL83/pKM005 (Fig. 1B, compare lanes 8 and 9 to lanes 2, 3, 5, and 6). These results indicate that the cellular adaptation to the low temperature is impaired with a more severe cold-shock response when cells harbor a multicopy plasmid carrying a part of the *cspA* gene.

Since the prolonged synthesis of *cspA* after cold shock was caused by pJG78, it was hypothesized that the 600-bp *cspA* upstream region cloned in pJG78 may sequester a factor  
10 responsible for the inhibition of *cspA* production after cold shock, resulting in the prolonged expression or the derepression of *cspA*. In order to examine this hypothesis, the 600-bp upstream region of *cspA* was re-cloned into pUC19. The plasmid is called pUC19-600. Note that the copy number of pUC19 (300 copies/cell) is about 10 times higher than pJG78 derived from pBR322 (30 copies/cell). A pulse-labeling experiment was carried out as  
15 described previously (Jiang *et al.*, 1993). As shown in Fig. 2, in the CL83 cells, CspA production increased up to 1.5 hr and was reduced to a basal level after 3 hr at 15°C (Fig. 2, lanes 1 to 6). In CL83 cells with pJG78, a certain level of *cspA* expression was still observed even after 24 hr at 15°C (Fig. 2, lanes 7 to 12). Patterns of *cspA* production in CL83 cells with pUC19-600 are similar to those with pJG78 (Fig. 2, lanes 13 to 18). However, the  
20 level of the *cspA* derepression was much higher with pUC19-600 than that with pJG78, as judged from the production of CspA at 3 and 5 hr. Thus, the higher the copy number of the *cspA* upstream region, the stronger the derepression of the *cspA* expression. Again, CsdA (indicated by X) showed the exactly same expression pattern as CspA throughout all the lanes shown in Fig. 2.

As shown in Fig. 1B, the cells with pJG78 showed a certain inhibition of general protein synthesis at low temperature (compare lanes 8 to 11 with lanes 2 to 5, respectively in Fig. 2). Significantly, this inhibition in the cells harboring pUC19-600 was even more evident than that in the cells harboring pJG78, in terms of both the protein synthesis rate and the inhibition time (compare lanes 14 to 17 with lanes 8 to 11, Fig. 2). The higher copy number of the 600-bp of *cspA* upstream region results in the stronger inhibition of the synthesis of other cellular proteins, indicating that cold-shock adaptation is inhibited.

#### EXAMPLE 4: Overproduction of the 5' untranslated region of the *cspA* mRNA.

In order to determine the precise region within the 600-bp sequence required for the derepression of *cspA* and the inhibition of cold-shock adaptation at low temperature, a series of internal fragments as shown in Fig. 3 were generated by PCR and cloned into the *Sma*I site of pUC19. Their sequences were confirmed by DNA sequencing. The ability to derepress expression of *cspA* and to inhibit cold-shock adaptation at 15°C for each construct was examined by pulse-labeling experiment. First, deletion mutations were made from the 5' end of the 600-bp fragment. As shown in Fig. 3, fragment 3 (186-base deletion), fragment 2 (312-base deletion), fragment 2E (366-base deletion) and fragment 2G (390-base deletion) all still retained the derepression function. Next, fragment 2 was further dissected into fragment 2A and 2B which overlap by 23 bp as shown in Fig. 3. Surprisingly, both 2A and 2B lost the functions. Fragment 2F which is longer by 33 bp at the 5' end than fragment 2B was also constructed, was still incapable of the functions. It was found here that the constructs which are capable of the derepression of *cspA* also result in inhibition of the cold-shock adaptation, and *vice versa*.

The fact that fragment 2 is functional for both the *cspA* derepression and the inhibition

of cold-shock adaptation, while fragment 2A is not, indicates that the *cspA* promoter region alone is not sufficient for the functions of the 600-bp fragment. Furthermore, the fact that functional fragment 2G is longer at the 5' end by 31 bp than the non-functional fragment 2F suggests a possibility that the both functions require the full *cspA* promoter for the transcription of the 5' UTR of the *cspA* mRNA. Note that the *cspA* mRNA has a 159-base untranslated sequence at the 5' end (Goldstein *et al.*, 1990). In order to confirm this possibility, the *cspA* transcripts produced from the cloned fragments (fragments 2, 2A, 2B, 2E, and 2F) were examined by primer extension. Using the total RNA fraction isolated from cells harboring various plasmids incubated for 1 hr at 15°C, primer extension was performed with two independent primers; primer 3550 which corresponds to the sequence from +124 to +143 in the 5' UTR and primer 3551 which corresponds to a part of the *cspA* coding sequence from +224 to +243. The former primer detects the *cspA* mRNA transcribed from both the plasmid and the chromosome, while the latter detects the mRNA only from the chromosomal *cspA* gene, since none of the plasmids contains the *cspA* coding region.

As shown in Fig. 4, the amounts of the transcript from the chromosomal *cspA* gene indicated by primer 3551 were basically the same among all constructs (Fig. 4, lanes 1 to 6). In contrast, the amount of the *cspA* transcripts encompassing the 5' UTR indicated by primer 3550 showed two different levels. For those nonfunctional constructs (pUC19-2A, pUC19-2B, and pUC-2F), the amounts of the transcripts detected by primer 3550 (lanes 3, 4, and 6 in Fig. 4, respectively) were almost identical to that with pUC19 (lane 1 in Fig. 4), indicating that the *cspA* regions cloned in these plasmids were not transcribed. On the other hand, for those functional constructs (pUC19-2 and pUC19-2E), much higher levels of the *cspA* transcripts detected by primer 3550 were observed (lanes 2 and 5 in Fig. 4, respectively) in comparison with the level with pUC19 (lane 1 in Fig. 4). These results demonstrate that the

5' UTR of the *cspA* mRNA was transcribed in fragment 2 and 2E, but not in fragments 2A, 2B and 2F. Therefore, the ability to prolong *cspA* expression and to inhibit the cold-shock adaptation at low temperature is clearly correlated with the transcription of the 5' UTR of the *cspA* mRNA.

5 In order to unambiguously demonstrate that the transcription of the 5' UTR of the *cspA* mRNA is required for both the *cspA* derepression and the inhibition of cold-shock adaptation, the entire promoter fragment (-457 to -1) plus 6-base (+1 to +6) region from *cspA* was cloned into pUC19. This fragment was designated fragment 1 (see Fig. 3). Thus, most of the 5' UTR of the *cspA* mRNA was deleted in fragment 1. By pulse-labeling experiment  
10 shown in Fig. 5B, fragment 1 was incapable of derepressing *cspA*, in spite of the fact that the transcripts from the *cspA* promoter were clearly detectable by primer extension (Fig. 5A). From these results, it is concluded that at least a portion of the *cspA* 5'UTR from +1 to +143 has to be transcribed to exert the effect on the *cspA* expression and the cold-shock adaptation.

15 **EXAMPLE 5: Cold-shock genes affected by the overproduction of the 5' UTR of the *cspA* mRNA.**

Next the overproduction of the 5' UTR of the *cspA* mRNA was examined to determine if the *cspA* mRNA has any effects on the expression of other cold-shock genes. The protein expression pattern of the cold-shocked cells overproducing the *cspA* 5' UTR was  
20 analyzed by two-dimensional electrophoresis. The plasmid pJYG21/X,S contains the entire *cspA* promoter and most of the 5' UTR of the *cspA* mRNA (+1 to +143), while pJYG81/X,S contains the entire *cspA* promoter but only the first 6-base region of the 5'UTR of the *cspA* mRNA. The cells harboring these plasmids were pulse-labeled as described before (Jiang *et al.*, 1993). At 37°C, the rate of protein synthesis and the protein pattern were very similar for

both strains (Fig. 6, A and B); note that no cold-shock proteins were detected. When these cells were shifted to 15°C for 1 hr (Fig. 6, C and D), the synthesis of cold-shock proteins (1) *cspA*; (2) *CspB'*; (3) *CspB*; and (4) *CsdA* became very prominent. Note that *CspB'* was co-induced with *CspB* and has been speculated to be either a modified form of *CspB* or a yet  
5 unidentified cold-shock protein (Etchegaray *et al.*, 1996). The rate of cold-shock protein synthesis for both constructs was comparable as judged from the densities of the spots. Although the synthesis of most other cellular proteins was significantly reduced for both strains compared with that at 37°C, much stronger inhibitory effects were observed in the cells transformed with pJG21/X,S. When cells were incubated at 15°C for 3 hr, synthesis of  
10 most cellular proteins recovered to a normal level with concomitant reduction of all the cold-shock proteins in the cells harboring pJG81/X,S (Fig. 6F). In contrast, for the cells harboring pJG21/X,S, the production of all the cold-shock proteins (marked by 1 to 4) was still maintained at a very high level along with reduced production of other cellular proteins (Fig. 6E). These results clearly demonstrated that overproduction of the 5' UTR of the *cspA*  
15 mRNA results in the derepression of not only *cspA* but also other cold-shock genes, suggesting that genes for cold-shock proteins are regulated by a common mechanism. It is also further confirmed that the inhibition of cold-shock adaptation is due to the overproduction of 5' UTR of the *cspA* mRNA by blocking the synthesis of other cellular proteins.

20 Based on the results described above, overproduction of the 5'UTR of the *cspA* mRNA causes the concomitant inhibition of other cellular proteins. This implies that cell growth upon cold shock would be more severely inhibited with the cells overproducing the 5'UTR of the *cspA* mRNA than that with the wild type cells. The growth of cells harboring pUC19-600 or pUC19-2G (see Fig. 3) was indeed severely inhibited. This was characterized



by a longer lag period (data not shown).

#### EXAMPLE 6: Effects of the overproduction of *cspA*

The hyper-expression of the 5' UTR of the *cspA* mRNA resulted in the prolonged overproduction of CspA (see Fig. 2). Therefore, the effects observed above may be due to the overproduction of the CspA protein rather than the 5' UTR of the *cspA* mRNA. This possibility was examined using CL83 cells harboring pJJG02 which contains the entire *cspA* gene. Pulse-labeling experiments were carried out as described above. As shown in Fig. 7, with strain CL83 carrying pUC19, the expression of *cspA* and *csdA* (the gene for protein X) were induced at 1 hr after the temperature shift to 15°C (lanes 1 and 2), and returned to a basal level at 3 hr after the temperature shift (lane 3). On the other hand, when the cells were transformed with pJJG02, the expression of *cspA* was not only induced at 15°C, but also significantly higher than that of cells with pUC19 as judged by two-dimensional gel electrophoresis (not shown). It should be noted that high CspA production is still observed even at 3 hr 15°C (lane 6). Although this overproduction of CspA at 3 hr after cold-shock was very similar to the case with the overproduction of the 5' UTR of *cspA* as described earlier (Fig. 2), it is important to note that no prolonged lag period of cell growth and no prolonged production of other cold-shock proteins such as CspB and CsdA were observed at the same time point. These results indicate that the co-production of CspA with the 5' UTR of the *cspA* mRNA suppresses the effects of the overproduction of only the 5' UTR, and that the high levels of CspA production even at 3 hr after cold-shock are not the cause of this effect.

#### EXAMPLE 7: Identification of a repressor binding site

We attempted to identify the specific region responsible for the *cspA* derepression within the 5' UTR of *cspA* mRNA. Because of the 11-base sequence commonly found in the 5' UTR in *cspA*, *cspB* and *cspA*, the region containing this sequence from position +1 to +25 of the 5'UTR of the *cspA* mRNA was tested. This region was put under the control of the *cspA* promoter in pUC19 to construct p2JTEK (Fig. 8). Cells harboring p2JTEK were then examined for the pattern of total protein synthesis at 3 hr after the temperature shift from 37 to 15°C. As shown in lane 3 in Fig. 8, CspA and CsdA production was clearly derepressed, with concomitant inhibition of other cellular proteins, as evident from the lighter background in comparison with cells harboring pUC19 (Fig. 8, lane 1). Cells harboring p6mTEK, which transcribes only the region from +1 to +6 of the *cspA* mRNA showed a pattern (Fig. 8, lane 2) identical to that of cells with pUC19 (Fig. 8, lane 1). This result indicates that the region responsible for the *cspA* derepression resides within the first 25-base sequence of the 5' UTR of *cspA* mRNA.

#### EXAMPLE 8: Deletion Analysis of the *cspA* 5'-UTR

Previously, we constructed two *cspA-lacZ* fusions in which the *lacZ* gene was transcriptionally fused to *cspA* at +26 (pKM67; Mitta *et al.*, 1997) or at +143 (pJG78; Jiang *et al.*, 1996) of the *cspA* mRNA. The  $\beta$ -galactosidase activity of the cells harboring pJG78 was very low at 37°C, and increased about 10 fold at 2 h after temperature downshift to 15°C, whereas the  $\beta$ -galactosidase activity of the cells harboring pKM67 was very high even at 37°C (Mitta *et al.*, 1997). It has been proposed that the region from +26 to +143 in the 5'-UTR of *cspA* mRNA has three possible functions on cold-shock induction: (i) repression of *cspA* expression at 37°C, (ii) stabilization of its mRNA upon cold shock, and (iii) translation efficiency at 15°C (Mitta *et al.*, 1997).

In order to further investigate which part of the 5'-UTR is responsible for the positive or negative regulation of *cspA* expression, we attempted deletion analysis of the 5'-UTR and examined the effect of various deletions on cold-shock induction of *cspA*. For this purpose, a series of 5'-UTR deletion mutants were constructed, in which a 26- to 32-base deletion was created at every about 30 bases, and the resultant 5'-UTR was translationally fused to *lacZ* at the 13th amino acid residue of CspA. The resultant plasmids, pMM022, pMM023, pMM024, pMM025 and pMM026 contain deletion mutations in the 5'-UTR from +2 to +27, from +28 to +55, from +56 to +86, from +86 to +117, and from +118 to +143, respectively (Fig. 9A). In the case of pMM024, a deletion from +56 to +85 was originally designed, but all the transformants we analyzed contained an extra base deletion at position +86. Plasmid pMM67 (Mitta *et al.*, 1997), which is the wild-type *cspA-lacZ* translational fusion construct, was used as a control. *E. coli* AR137, a *pcnB* mutant, which is known to maintain pBR322 derivatives in a low copy number (Lopilato *et al.*, 1986), was used as a host for transformation in order to avoid multicopy effects of the constructed gene on their expression.

Transformed cells were grown in M9-Casamino acids medium at 37°C, and  $\beta$ -galactosidase activities were measured after temperature downshift from 37°C to 15°C. At 37°C (zero time point in Fig. 9B),  $\beta$ -galactosidase activities were 10 fold higher in cells harboring pMM024 ( $\Delta$ 56-86) and pMM025 ( $\Delta$ 86-117) than in cells harboring the wild-type pMM67, while other deletion mutants [pMM022 ( $\Delta$ 2-27), pMM023 ( $\Delta$ 28-55) and pMM026 ( $\Delta$ 118-143)] showed very low  $\beta$ -galactosidase activities. These results suggest that the 5'-UTR region from base +56 to +117 is involved in the repression of *cspA* expression at 37°C. Interestingly  $\beta$ -galactosidase activity increased almost 5 fold with pMM024 ( $\Delta$ 56-86) after temperature downshift, while it increased only less than 2 fold with pMM025 ( $\Delta$ 86-117), suggesting that the region deleted in pMM025 ( $\Delta$ 86-117) plays an important role in cold-

shock induction of *cspA*. Similar to pMM025 ( $\Delta 86-117$ ),  $\beta$ -galactosidase activity with pMM023 ( $\Delta 28-55$ ) was poorly induced at low temperature. In particular, the region deleted in pMM026 ( $\Delta 118-143$ ) appears to play a crucial role in *cspA* expression at both high and low temperatures, since  $\beta$ -galactosidase activity was very low at both 37°C and 15°C (Fig. 9B). The deletion of the region from base +2 to +27 (pMM022) containing the cold-box sequence involved in *cspA* autoregulation (Jiang *et al.*, 1996) has little effect on the cold-shock induction of *cspA* as predicted (Fig. 9B).

#### EXAMPLE 9: Analyses of *cspA-lacZ* mRNA

As described above, all deletion mutations of the 5'-UTR except for pMM022 ( $\Delta 2-27$ ) affected on cold-shock induction of *cspA*. The *cspA* promoter is known to be active even at 37°C (Goldenberg *et al.*, 1997; Fang *et al.*, 1997; Mitta *et al.*, 1997). Since all the deletion constructs have the intact *cspA* promoter (Fig. 9A), transcription efficiencies of these constructs are likely to be identical. On the other hand, the *cspA* mRNA stability is significantly different depending on growth temperatures (Brandi *et al.*, 1996; Goldenberg *et al.*, 1996; Bae *et al.*, 1997; Fang *et al.*, 1997; Goldenberg *et al.*, 1997; Mitta *et al.*, 1997). Therefore the effect of the deletion mutations on the *cspA* expression at low temperature may be due to different mRNA stabilities of the constructs. To examine this aspect, the primer extension analysis was carried out to quantitate the amounts of the *cspA-lacZ* transcripts for each mutant at different time points after the addition of rifampicin at both 37°C and 15°C (Fig. 10A). Again, in order to avoid multicopy effects either positively or negatively, strain AR137, a *pcnB* mutant, was used. The amounts of transcripts at each time point were estimated by a phosphorimager, and the amounts of mRNA remained (per cent of the amount at zero time point) were plotted as shown in Fig. 10B. All the transcripts were unstable at

37°C with their half-lives estimated between 30 and 45 s. At 15°C, however, they became very stable with half-lives between 20 and 40 min. These half-lives are similar to those for the wild-type construct pMM67 obtained previously (Mitta *et al.*, 1997) as well as to those for the wild-type chromosomal *cspA* (Fang *et al.*, 1997; Goldenberg *et al.*, 1997). It is important to note that in contrast to the similar mRNA half-lives at low temperature,  $\beta$ -galactosidase activities induced at 15°C were widely varied among all these constructs as shown in Fig. 9B.

These results indicate that *cspA* induction efficiencies as measured by  $\beta$ -galactosidase activity are not correlated to the mRNA stability of each deletion construct, but rather to the amount of mRNA and/or its translation efficiency for each construct. Therefore, we next examined the amounts of mRNA for each construct at different time points after cold shock by the primer extension method. The results are shown in Fig. 11A and the amounts of transcripts were estimated by a phosphorimager. Their relative amounts were calculated using the amount of the transcript of pMM67 at zero time as 1 (Fig. 11B). At 37°C, the amounts of transcripts for pMM022 ( $\Delta 2-27$ ) and pMM024 ( $\Delta 56-86$ ) were very similar to that of the wild-type construct pMM67 (column 1 in Fig. 11B). It should be noted that  $\beta$ -galactosidase activity of pMM024 ( $\Delta 56-86$ ) at 37°C was more than 10 times higher than that of pMM022 ( $\Delta 2-27$ ) (see Fig. 9B). In the case of pMM023 ( $\Delta 28-55$ ), pMM025 ( $\Delta 86-117$ ) and pMM026 ( $\Delta 118-143$ ), the amounts of transcripts at 37°C are approximately half of that of pMM67. Again it should be noted that  $\beta$ -galactosidase activity of pMM025 ( $\Delta 86-117$ ) was 10 times higher than those of pMM023 ( $\Delta 28-55$ ) and pMM026 ( $\Delta 118-143$ ) (Fig. 9B). These results indicate that there is no correlation between the amounts of transcripts and the  $\beta$ -galactosidase activities at 37°C.

#### EXAMPLE 10: Translational Regulation by the 5'-UTR

After temperature downshift, the amounts of the *cspA-lacZ* mRNAs dramatically increased in all the constructs and the induction patterns are shown in Fig. 11A and 11B. They showed very similar pattern in accumulation of the transcripts as that of the wild-type pMM67, such that the maximal induction was observed at 1 h after temperature downshift.

5 The patterns of mRNA levels were very similar between pMM67 and pMM024 ( $\Delta 56-86$ ), while the others also showed a similar induction pattern although the amounts of their mRNAs were approximately a half of that of the pMM67 mRNA at each time point. Since the promoter activity of all the deletion constructs are considered to be the same, and in addition their mRNA stabilities were also very similar to that of the wild-type construct (Fig.

10 10B), lower amounts of mRNAs for all the deletion constructs except for pMM024 ( $\Delta 56-86$ ) are probably due to their slower transcription elongation rate and/or transcription attenuation within the 5'-UTR. A remarkable finding was that the amounts of mRNA for pMM026 ( $\Delta 118-143$ ) accumulated after cold shock were also identical to those for pMM022 ( $\Delta 2-27$ ), pMM023 ( $\Delta 28-55$ ) and pMM025 ( $\Delta 86-117$ ) (see Fig. 11B). Nevertheless cold-shock

15 induction of  $\beta$ -galactosidase activity was extremely low for pMM026 ( $\Delta 118-143$ ) throughout cold-shock treatment (Fig. 9B), indicating that the mRNA for this construct was very poorly translated. Relative translation efficiencies at 15°C was calculated for all the constructs from the increments of  $\beta$ -galactosidase activity during the cold shock and the amounts of mRNA (see the Brief Description of the Figures Legend to Fig. 11). As shown in Fig. 11C, the

20 translation efficiency of pMM026 ( $\Delta 118-143$ ) mRNA was extremely poor and calculated to be 9.5% of that of pMM67 mRNA (Fig. 11C column 6). Thus, it is clear that the deletion from base +118 to +143 of the 5'-UTR negatively affected on translation. Besides pMM026 ( $\Delta 118-143$ ), the translation efficiencies of the mRNAs of pMM023 ( $\Delta 28-55$ ) and pMM025 ( $\Delta 86-117$ ) were relatively low and calculated to be 54 and 36% of that of pMM67 mRNA,

respectively. These results clearly indicate that the translation efficiency of mRNA plays a crucial role in the regulation of *cspA* expression and that in particular, the region from base +118 to +143 of the 5'-UTR plays a major role in translation efficiency.

Nucleotide sequence comparison of the 5'-UTR regions of the cold-shock inducible genes, *cspA*, *cspB* and *cspG*, reveals that there is a 13-base sequence (from base +123 to +135 of *cspA*) conserved very well among these genes as shown in Fig. 12. Interestingly, these 13-base sequences are located immediately upstream of the Shine-Dalgarno sequence and contains a palindromic sequence to form a stable secondary structure ( $\Delta G = -9.5\text{kcal}$ ; see Figs. 12 and 14). It is also complementary to the region from base 1023 to 1035 of 16S rRNA (Fig. 12). In the pMM026 ( $\Delta 118-143$ ) mRNA, this upstream-box region is deleted, which may be the major cause for the poor translation efficiency of the mRNA.

In order to characterize the role of the upstream box in translation efficiency, two new constructs were made; in one construct (pKNJ37) the exact 13-base upstream-box sequence was deleted from the wild-type *cspA-lacZ* construct (pMM007) and in another (pKNJ38) the 13-base sequence was added at the upstream region of SD of pKM67 (Mitta *et al.*, 1997) as shown in Fig. 13A. In pKM67 the *lacZ* gene is fused at base +26, and it has been shown that as a result of the substantial deletion in the 5'UTR  $\beta$ -galactosidase became expressed even at 37°C without any further induction upon cold shock (Mitta *et al.*, 1997). Note that both pMM007 and pMM67 have the exactly identical insert of *cspA* in different vectors, pRS414 and pKM005, respectively. Cold-shock induction patterns of  $\beta$ -galactosidase activities of these two plasmids (pMM007 and pMM67) are similar (data not shown). Cells were transformed with pKNJ37, pMM007, pKNJ38 and pKM67, and cold-shock induction of  $\beta$ -galactosidase was examined. Deletion of the upstream box (pKNJ37) significantly lowered  $\beta$ -galactosidase activity not only at 37°C but also upon cold shock (Fig.13B). Contrary to the

result, when the upstream box was inserted into pKM67 13 bases upstream of the SD sequence, the constitutive expression of  $\beta$ -galactosidase at 37°C increased by approximately 20% (Fig.13B). This increment was also kept upon cold shock. These results support the notion that the upstream-box sequence is associated with efficient translation of *cspA*.

5 Specific sequences of the 5' UTR which are useful in the invention include 5'-GCCGAAAGGCACA-3' (SEQ ID NO:48), 5'-GCCGAAAGGCUCA-3' (SEQ ID NO:49), and 5'-GCCGAAAGGCCCA-3' (SEQ ID NO:50). (See Fig. 12).

It is currently considered that *cspA* expression is regulated at the levels of transcription, mRNA stability and translation as follows: (i) The *cspA* gene has a strong promoter equipped with the UP element, which works at both 37°C and 15°C (Fang *et al.*, 1997; Goldenberg *et al.*, 1997; Mitta *et al.*, 1997), although transcription of the *cspA* gene does not require any *de novo* protein synthesis upon cold shock. (ii) The *cspA* mRNA contains the downstream-box (DB) sequence downstream of the initiation codon, which plays a major role in enhancement of translation initiation at low temperatures (Etchegaray and Inouye, unpublished; Mitta *et al.*, 1997). (iii) It is very important to note that the *cspA* mRNA has an unusually long 5'-UTR (Tanabe *et al.*, 1992), consisting of 159 bases, and is extremely unstable at 37°C (Brandi *et al.*, 1996; Goldenberg *et al.*, 1996; Bae *et al.*, 1997; Fang *et al.*, 1997; Jiang *et al.*, 1997; Goldenberg *et al.*, 1997; Mitta *et al.*, 1997). Immediately upon cold shock, the *cspA* mRNA becomes stable. Again, this stabilization of mRNA upon cold shock does not require any *de novo* protein synthesis. When the region from base +26 to +143 of the 5'-UTR was deleted from the *cspA-lacZ* fusion construct, high  $\beta$ -galactosidase activity was obtained even at 37°C (Mitta *et al.*, 1997). In the 5'-UTR, there is a putative RNaseE cleavage site immediately upstream of the SD sequence (Fang *et al.*, 1997). This site is considered to be responsible for the extreme instability of the *cspA* mRNA, since the three-



base substitution mutation at this region resulted in 150-fold stabilization of the mRNA, allowing a high CspA production even at 37°C (Fang *et al.*, 1997). Thus, it is clear that the unusually long 5'-UTR of the *cspA* mRNA is responsible for its instability at 37°C. It can be considered that the 5'-UTR makes the *cspA* mRNA extremely unstable at 37°C, in such a way as *cspA* is cold-shock inducible. It is worth mentioning that although the *cspA* mRNA becomes stable and is accumulated at the nonpermissive temperature in the temperature-sensitive RNaseE mutant, CspA production was not detected under this condition (Fang *et al.*, 1997), suggesting that in addition to the stability of mRNA another role may exist in the 5'-UTR.

We have attempted to further elucidate the roles of the unusually long 5'-UTR of the *cspA* mRNA in *cspA* expression. We made a series of 26- to 32-base deletion mutations encompassing the entire 5'-UTR. These mutated 5'-UTR regions were translationally fused to *lacZ* at the 13th amino acid residue of CspA after the DB sequence. At 37°C, pMM022 ( $\Delta$ 2-27), pMM023 ( $\Delta$ 28-55) and pMM026 ( $\Delta$ 118-143) showed the similar  $\beta$ -galactosidase activities as the wild-type pMM67, while pMM024 ( $\Delta$ 56-86) and pMM025 ( $\Delta$ 86-117) showed more than 10 fold higher  $\beta$ -galactosidase activities than pMM67, indicating that the region from base +56 to +117 is involved in the repression of *cspA* expression at 37°C.

Based on the  $\beta$ -galactosidase activities after temperature downshift, mutants can be classified into three classes; those in Class I [pMM022 ( $\Delta$ 2-27) and pMM024 ( $\Delta$ 56-86)] in which  $\beta$ -galactosidase is induced in a similar fashion to pMM67, those in Class II [pMM023 ( $\Delta$ 28-55) and pMM025 ( $\Delta$ 86-117)] in which cold-shock induction of  $\beta$ -galactosidase is poor, and that in Class III [pMM026 ( $\Delta$ 118-143)] with very low  $\beta$ -galactosidase activities both before and after cold shock. It is worth mentioning that these differences were due to neither the amounts nor the stability of mRNA as evident from Figs. 10 and 11. This supports the

notion that the 5'-UTR has another role in translation efficiency in addition to the stability of mRNA as mentioned above.

The relative translation efficiencies after temperature downshift for different constructs showed surprisingly significant differences (Fig. 11C), which coincide well with the classification of the constructs at 15°C as described above. The translation efficiencies with pMM022 ( $\Delta$ 2-27) and pMM024 ( $\Delta$ 56-86) (Class I) are a little better than that of the wild-type pMM67, those with pMM023 ( $\Delta$ 28-55) and pMM025 ( $\Delta$ 86-117) (Class II) are 40 to 50% of that of pMM67, and that with pMM026 ( $\Delta$ 118-143) (Class III) is less than 10% of that of pMM67.

In pMM026 ( $\Delta$ 118-143), the deletion mutation is clearly affecting the translation efficiency but not the stability of mRNA. The deleted region was found to contain a 13-base sequence (base +123 to +135) well conserved in the mRNAs for all the cold-shock inducible *cspA* family, *cspA*, *cspB*, *cspG*, and *cspI* (see Fig. 12). This sequence designated the upstream box may form a distinct secondary structure in both the wild-type pMM67 and Class I constructs [pMM022 ( $\Delta$ 2-27) and pMM024 ( $\Delta$ 56-86)] (Fig. 14). Class I constructs showed a similar translation efficiency to the wild-type. In Class II [pMM023 ( $\Delta$ 28-55) and pMM025 ( $\Delta$ 86-117)], which showed 50% of translation efficiency of the wild-type, this secondary structure disappears, however, the predicted secondary structures around the SD sequence in these constructs are still similar to that of the wild-type construct. In contrast, when the upstream-box region is deleted [pMM026 ( $\Delta$ 118-143); Class III, which showed a very poor translation efficiency], the SD region forms a more stable secondary structure. This likely prevents recognition of the mRNA by ribosomes, causing a very poor translation efficiency in pMM026 ( $\Delta$ 118-143). Therefore the upstream box may function to punctuate the formation of a stable secondary structure immediately upstream of the SD sequence,

allowing it highly accessible to ribosomes. Alternatively, as the upstream-box sequence is complementary to the 16S rRNA sequence from base 1023 to 1035 (see Fig. 12), it is possible that the upstream-box sequence may be another *cis*-element, which may enhance translation efficiency by forming a duplex with 16S rRNA in addition to the SD sequence and the downstream box (Mitta *et al.* 1997). Since all the constructs use the identical site of *cspA* to fuse to *lacZ*, the observed differences in translation efficiency are considered to be at the level of translation initiation but not at the level of translation elongation. It has been suggested that such an interaction between mRNA and 16S rRNA plays an important role in translation initiation (McCarthy and Brimacombe, 1994). The region from base 1023 to 1035 of 16S rRNA has been shown to be near the site where 30S ribosomes interact with mRNA on the basis of the fact that the U residue at position 1052 directly interacts with the 6th base from the initiation codon (Dontsova *et al.*, 1992; Rinke-Appel *et al.*, 1993). Furthermore, it has been reported that RNase V1 sensitivity at G1020, A1021 and A1022 increased immediately after the assembly of 30S ribosomal subunits, indicating that the site encompassing these residues is exposed to the surface (Powers *et al.*, 1988).

Consistent with the proposed role of the upstream box in the translation, the addition of an upstream-box sequence to pKM67 resulted in increase of  $\beta$ -galactosidase activity approximately by 20%. On the other hand, the deletion of the exact 13-base upstream-box sequence [pKNJ37 ( $\Delta$ 123-135) in Fig. 13] resulted in 50% reduction of  $\beta$ -galactosidase activity at 37°C and a lower level of induction upon cold shock. Note that the predicted secondary structure surrounding the SD sequence of pKNJ37 ( $\Delta$ 123-135) is the same as the wild-type (Fig. 14).

In the heat shock response, the synthesis of the heat-shock sigma factor,  $\sigma^{32}$ , is regulated at the level of translation and the secondary structure of the *rpoH* mRNA plays a

crucial role in this regulation (Nagi *et al.*, 1991; Yuzawa *et al.*, 1993). Very recently, the secondary structure of the *rpoH* mRNA has been determined by chemical and enzymatic probing assays and the results are completely consistent with the predicted *rpoH* secondary structure proposed previously (Morita *et al.*, 1999). It has also been shown that mutations in the *rpoH* mRNA, which are predicted to decrease the mRNA stability, increased *rpoH* expression and *vice versa* (Morita *et al.*, 1999). Toeprinting assays using the wild-type *rpoH* mRNA and 30S ribosomes showed that the toeprinting appearance totally depends on temperature, indicating that the *rpoH* mRNA secondary structure at 42°C, which is accessible to ribosomes, is different from that at 30°C, and that the *rpoH* mRNA structure itself is a determinant for *rpoH* expression without any other factors (Morita *et al.*, 1999). Thus, it was proposed that the *rpoH* mRNA acts as an RNA thermosensor (Morita *et al.*, 1999). Storz (1999) discussed that the *lcrF* mRNA of *Yersinia pestis* (Hoe *et al.*, 1993) and the  $\lambda$  phage cIII mRNA (Altuvia *et al.*, 1989) might be other examples for RNA thermosensor.

In summary, stabilization of the *cspA* mRNA upon cold shock is prerequisite for CspA production. This stabilization does not require any *de novo* protein synthesis. As presented here, translation efficiency, which may involve the secondary structure of the 5'-UTR, in particular surrounding the SD sequence, of *cspA* mRNA, is turned out to play an important role in *cspA* expression in addition to the mRNA stabilization. It is thus possible that the *cspA* mRNA might act as an RNA thermometer as proposed for the *rpoH* mRNA. To know the more precise molecular mechanism of the regulation of *cspA* expression, determination of the secondary structure of the *cspA* mRNA and the relationship between the secondary structure and the translation efficiency remain to be addressed. The point mutation analysis in addition to the deletion analysis presented here will give us information on the molecular anatomy of the structure and function of 5'-UTR of the *cspA* mRNA.

**Example 11: The Role of DB in the Cold-shock Induction of *cspB*, and the Effect of a Perfectly Matching DB**

Previously, we have demonstrated that the expression of *cspB* at low temperature is primarily regulated at the mRNA level (Etchegaray et al., 1996). Here, we first show that DB is an important contributor to the translational induction of *cspB* at low temperature. *E. coli* AR137, a *pcnB* mutant which maintains pBR322 derivative plasmids at low copy (Lopilato et al., 1986) was transformed with a series of *cspB-lacZ* translational fusions (Fig. 15A and 15B). Figure 15C shows  $\beta$ -galactosidase activity at various time points after temperature shift from 37 to 15°C. After 3 hr at 15°C, pB13 and pB17 show the highest levels of  $\beta$ -galactosidase activity, indicating that a region from codon 3 and 13 (containing DB) plays a major role in high expression of *cspB* at low temperature. Levels of  $\beta$ -galactosidase units from the SD-deletion constructs pB13sd and pB17sd demonstrate that SD is also required for the induction of *cspB*. Primer extension analysis (Fig. 15D) shows that the mRNA levels of pB3 and pB17 are almost identical, while the  $\beta$ -galactosidase activity of pB17 is 7 times higher than that of pB3 (Fig. 15C), demonstrating that the differences in the sequence downstream of the initiation codon cause a significant effect on the efficacy of mRNA translation but not on the amount of the mRNAs. Based on the amounts of mRNA estimated using a phosphorimager and the increments of  $\beta$ -galactosidase activity between 1 and 2 hr. after cold-shock induction, the translational capability of pB17 was calculated to be 6 times higher than that of pB13 and 18 times higher than that of pB3. This could be explained by the potential to form additional base-pairing between the *cspB*-mRNA (pB17) and the 16S rRNA as shown in Figure 15A. This suggests that a longer DB (pB17) could be more effective for translation than a shorter DB (pB13). When the SD sequence was deleted from

pB13, the translational efficiency became even lower than that of pB3, indicating that both SD and DB are required for the full expression of *cspB* at low temperature.

It has been shown that mRNA stability plays a role in the cold-shock inducibility of *cspA* (Brandi *et al.*, 1996; Goldenberg *et al.*, 1997; Fang *et al.*, 1997). However, the dramatic effect of DB on the *lacZ* expression cannot account for the differences of the mRNA stabilities at 15°C between the constructs with and without DB. Fig. 15E shows the half-life for pB3, pB13, pB13sd and pB17 to be 12, 22, 15 and 20 minutes, respectively.

It has been shown that mRNA stability plays a role in the cold-shock inducibility of *cspA* (Brandi *et al.*, 1996; Goldenberg *et al.*, 1997; Fang *et al.*, 1997). However, the dramatic effect of DB on the *lacZ* expression cannot account for the differences of the mRNA stabilities at 15°C between the constructs with and without DB. Figure 1E shows the half life for pB3, pB13, pB13sd, and pB17 to be 12, 22, 15 and 20 minutes, respectively.

It has been previously shown that DB is essential for the production of CspA at low temperature (Mitta *et al.*, 1997). However, the wild-type DB of *cspA* has 10 matches out of 15 possible matches to the anti-DB in 16S rRNA (Mitta *et al.*, 1997). Therefore, we added DBs of 12 (pJG78DB1) or 15 (pJG78DB2) bases that are complementary with the anti-DB of 16S rRNA to the site after the 5th codon of *lacZ* under the *cspA* regulatory system in pJG78 (see Fig. 15A) to examine if they enhance *lacZ* expression at 15°C. Mid-log phase cells (*penB*) grown at 37°C were shifted to 15°C and  $\beta$ -galactosidase activity was measured at 1, 2 and 3 hr after the shift. Fig. 15B shows that at 1 hr at 15°C the  $\beta$ -galactosidase activity was 3 and 8 fold higher with pJG78DB1 and pJG78DB2, respectively than with pJG78. After 2 and 3 hr at 15°C, the  $\beta$ -galactosidase activity was increased 3.5 and 10.5 times with pJG78DB1 and pJG78DB2, respectively than with pJG78. Moreover, the effect of the DB was observed at 37°C in which the  $\beta$ -galactosidase activity of pJG78DB1 and pJG78DB2

was 2 and 4 fold higher as compared with pJG78. The amount of the *lacZ* mRNA (Fig. 15C) as well as the mRNA stability did not vary significantly between these constructs. The *lacZ* mRNA half-life from pJG78, pJG78DB1 and pJG78DB2 was calculated to be 27, 23 and 25 min respectively. In addition, computer analysis (Zuker and Stieger 1982) revealed no significant differences in the mRNA secondary structures among pJG78, pJG78DB1 and pJG78DB2, suggesting that the insertion of the perfectly matching DB may not have a particular effect in the mRNA secondary structures that could account for the difference in their  $\beta$ -galactosidase expression. These results indicate that DB function as a translational enhancer and that greater complementarity to the anti-DB improves translational efficiency and/or that specific base-pairings like the first 3 nucleotides of the DB from pJG78DB2 may play an important role for the DB activity.

#### Example 12: Analysis of DB Function

The experiments described above were carried out at 15°C. In order to examine whether DB also works at 37°C, the *cspA* cold-shock regulatory regions upstream of SD of pJG78 and pJG78DB2 were replaced with the constitutive *lpp* promoter and the *lac* promoter-operator fragment using a pINIII vector (Inouye, M. 1983), yielding pINZ and pINZDB1, respectively (Fig. 17A). Cells (*pcnB*) transformed with pINZ or pINZDB1 showed very low  $\beta$ -galactosidase activity in the absence of IPTG, an inducer of the *lac* promoter (Fig. 1C, time 0). Upon the addition of 1mM IPTG,  $\beta$ -galactosidase activity was induced in both cells. After 3 hr induction,  $\beta$ -galactosidase activity increased 18 and 37 folds for pINZ and pINZDB1, respectively (Fig. 17C). However, the levels of  $\beta$ -galactosidase activity show a dramatic difference between the two; the activity with DB (pINZDB1) was 34 times higher than that without DB (pINZ), demonstrating that DB functions at 37°C as well.

Specific activities of  $\beta$ -galactosidase produced from vector pINZ and pINZDB1 are almost identical (data not shown) and thus the addition of the 5 amino acid residues in the  $\beta$ -galactosidase sequence of pINZDB1 (due to DB) does not affect the enzymatic activity. Furthermore, the stabilities of  $\beta$ -galactosidase from pINZ and pINZDB1 are also identical with a half-life of approximately 3 hr (data not shown).

Next, we examined whether DB functions independently from SD, the initial ribosomebinding site. For this purpose, the SD sequence of pINZDB1, GAGG was changed to GCCC, yielding pINZDB2 (Figure 17A and 17B). The  $\beta$ -galactosidase level of pINZDB2 induction was reduced to 1/7 of that of pINZDB1, but still 3 fold higher than that of pINZ at 10 hr after IPTG induction (Figure 17C). However, since pINZDB2 has the second AUG codon 6 codons downstream as a result of DB insertion it might serve as a secondary initiation codon for the *lacZ* gene. Indeed, the N-terminal sequence analysis showed that 100% of the  $\beta$ -galactosidase produced from pINZDB2 is initiated at the second AUG codon as compared with pINZDB1, which is more than 90% initiated at the first AUG codon (data not shown). Furthermore, the second AUG codon is preceded by a potential but poor SD sequence (AAGG) at the region corresponding to the 2nd and 3th codons (underlined; Fig. 17B). Indeed, when this secondary SD was removed by deletion of the 15-base sequence (codon 1 to 5; pINZDB3 in Fig. 17B),  $\beta$ -galactosidase activity at all time points was reduced to the background level (Fig. 17C), indicating that the secondary SD played a crucial role in the translation of the pINZDB2 *lacZ* mRNA. When the SD sequence was recreated by 5 base substitution in pINZDB3 (pINZDB4; Fig. 17B),  $\beta$ -galactosidase activity of this construct was recovered to a comparable level to that of pINZDB1 (Fig. 17C). It is important to notice that the DB sequence starting from the first AUG codon was eliminated in pINZDB4 (Fig. 17B). Therefore, the high expression of  $\beta$ -galactosidase from pINZDB1 and pINZDB4 is due to the



perfectly matching DB sequence (Fig. 17B). These results indicate that (a) DB functions only in the presence of SD, (b) the position of DB is flexible starting from either codon 1 or 6.

### Example 13: Enhancement of Translation by DB

5           The  $\beta$ -galactosidase activity showed in Fig. 17C indicates that DB enhances the translation of pINZDB1. Therefore, in order to test the effect the DB in translation efficiency the rate of  $\beta$ -galactosidase synthesis from pINZ and pINZDB1 was analyzed. The rate of  $\beta$ -galactosidase synthesis was measured by pulse-labeling cells for 5 min with [ $^{35}$ S]-methionine after the addition of IPTG using cells harboring pINZ and pINZDB1. After SDS-PAGE, the  
10           amounts of radioactive  $\beta$ -galactosidase were estimated using a phosphorimager (Fig. 17D). Prior to the addition of IPTG, the rate of  $\beta$ -galactosidase synthesis from pINZ and pINZDB1 was identical. However, upon IPTG induction the rates of  $\beta$ -galactosidase synthesis from pINZDB1 was continuously increasing at each time point while the rate of  $\beta$ -galactosidase synthesis from pINZ was almost non affected. After 4 hr of IPTG addition the rate of  $\beta$ -  
15           galactosidase synthesis from pINZDB1 was 6.5 times higher than that of pINZ. This result demonstrate that DB enhances the translation efficiency of pINZDB1 as reflected by the increment in the synthesis of  $\beta$ -galactosidase.

          In order to examine whether DB enhances translation initiation we next analyze the ability of *lacZ* mRNA from pINZ and pINZDB1 to form polysomes. For this experiment,  
20           *pcnB*<sup>+</sup> cells were used to amplify the effect of DB. Interestingly, cells with pINZDB1 could not form colonies on LB plates in the presence of 1 mM IPTG, while cells with pINZ formed colonies. The lethal effect of IPTG on the cells with pINZDB1 is considered to be due to overexpression of  $\beta$ -galactosidase. After the addition of IPTG, cell growth was stopped by the addition of chloramphenicol (0.1 mg/ml) at 15, 30 and 60 min and then polysome profiles

were examined as shown in Fig. 17D. From each gradient fraction (500 ml), 200 ml were spotted on a nitrocellulose membrane and the amount of the *lacZ* mRNA analyzed using a 24-base antisense oligonucleotide (M13-47 oligonucleotide). The amounts of the *lacZ* mRNA were quantified by a phosphorimager and are displayed in Fig. 17D. While the polysome profiles are similar, there are significant differences in the distribution of the *lacZ* mRNA; at 15 min the *lacZ* mRNA mainly exists in the upper half of the gradient (fraction 8 to 14, corresponding to 70S to 30S ribosomes) with pINZ, while with pINZDB1 a major peak (fraction 3 to 8) is formed in the lower half of the gradient. At 30 min, the *lacZ* mRNA from pINZ moved to the position of 70S ribosome, while the *lacZ* mRNA from pINZDB1 maintained a similar pattern as that at 15 min. At 60 min a major fraction of the *lacZ* mRNA from pINZ remained in the upper half of the gradient, while the *lacZ* mRNA from pINZDB1 was broadly distributed from higher order polysomes to 70S ribosome fraction. Therefore, the reason why cells harboring pINZDB1 could not form colonies on LB plates containing 1 mM IPTG may be due to a decrease in the concentration of free ribosomes as a result of the massive expression of a highly translatable DB-containing mRNA (Vind *et al.*, 1993). These results indicate that DB enhances the efficiency of polysome formation probably due to a translation initiation enhancement.

In order to estimate the exact effect of DB from the above experiment, the amount of the *lacZ* mRNA and the  $\beta$ -galactosidase activity were measured at the same time points taken in the polysome profiles (15, 30 and 60 min after IPTG induction). As shown in Fig. 18A, the amounts of the *lacZ* mRNA reached almost the maximal level at 15 min for both pINZ and pINZDB1. The phosphorimager analysis of this result revealed that the amounts of the pINZDB1 mRNA are 1.5, 1.4 and 1.3 times higher than those of the pINZ mRNA at 15, 30 and 60 min, respectively. The higher mRNA levels for pINZDB1 are probably attributable to

the highly efficient polysome formation of pINZDB1 that may stabilize the mRNA (Iost and Dreyfus 1995). The induction of  $\beta$ -galactosidase activity is shown in Fig. 18B. In the case of pINZDB1, the activity is very high even in the absence of IPTG, and upon the addition of IPTG, it increased from 18,500 to 64,400 units (3.5 fold) after 2.5 hr incubation. In the case of pINZ, the background activity prior to IPTG induction was much lower, and it increased from 900 to 2,900 units (3 fold) at the 2.5 hr time point. The increment of the  $\beta$ -galactosidase activity of pINZDB1 between 30 and 60 min is 35 times higher than that of pINZ, and therefore the efficiency of  $\beta$ -galactosidase production for pINZDB1 is calculated to be 26 times higher than that for pINZ on the bases of the amount of mRNA. Therefore, the higher levels of  $\beta$ -galactosidase production from pINZDB1 are due to a high efficiency of polysome formation.

Next, in order to more directly demonstrate the translation-enhancement effect of DB the  $\beta$ -galactosidase synthesis was examined in a cell-free system using pINZ and pINZDB1. The [ $^{35}$ S]methionine incorporation into  $\beta$ -galactosidase (band G) with pINZDB1 (lane 2, Fig. 19) was 8 fold higher than that with pINZ (lane 1), while the  $\beta$ -lactamase (band L) production was almost identical in both lanes.

It has been proposed that in the absence of ribosomal protein S2, structural changes in 16S rRNA result in the release of the anti-DB sequence from the penultimate stem making it more accessible to base-pair with DB (Shean and Gottesman 1992; Powers *et al.*, 1988). We analyzed the  $\beta$ -galactosidase expression of pINZ, pINZDB1 in *E. coli* CS239 that carries an S2 temperature-sensitive mutation (Shean and Gottesman, 1992). Fig. 20A shows that the  $\beta$ -galactosidase activity of pINZDB1 significantly increases upon shifting the temperature from 30 to 42°C in the S2<sup>ts</sup> strain (CS239) (6.3 fold from 0 to 3.5 hr), while the activity in the wild type strain (CS240) slightly increased (1.1 fold from 0 to 3.5 hr). If the initial ratio of the

activity of CS239 to that of CS240 at time 0 is taken as one, the ratio dramatically increased, reaching 5.8 at 3.5 hr after temperature shift (Fig. 20B). In contrast, the *lacZ* gene without DB did not show any significant differences in its expression between CS240 and CS239, and the ratios of the activity of CS239 to that of CS240 remained also at the initial level throughout the incubation time (Fig. 20B). A similar experiment was carried out with pINZDB3 (SD<sup>-</sup>, DB<sup>+</sup>), and the ratio of the activity in CS239 to that in CS240 increased 3.4 fold at 3.5 hr after the temperature shift (data not shown). These results clearly demonstrate that the low levels of S2 protein at 42°C causes significant stimulation of the *lacZ* expression only if the *lacZ* gene contains DB, consistent to the proposal of Shean and Gottesman (1992).

## REFERENCES

1. Sprengart, M. L., Fatscher, H. P. And Fuchs, E. (1990). Nucleic acids Res. 18, 1719-1723.
2. Sprengart, M. L., Fuchs, E. and Porter, A. G. (1996). EMBO J. 15, 665-674
3. Shean, C. S., and Gottesman, M. E. (1992). Cell 70, 513-522
4. Powers, T., Stern, S., Changchien, L. and Noller, H. F. (1988). J. Mol. Biol. 201, 697-716
5. Resch, A., tedin, K., Grundling, A., Mundlein, A., and Blasi, U. (1996). EMBO J. 15, 4740-4748
6. Sprengart, M. L. and Porter, A. G. (1997). Mol. Microbiol. 24, 19-28
7. Mitta, M., Fang, L., and Inouye, M. (1997). Mol. Microbiol. 26, 321-335
8. Ringquist, S., Jones, T., Snyder, E. E., Gibson, T., Boni, I., and Gold, L. (1995). Biochemistry 34, 3640-3648
9. Etchegaray, J-P., Xia, B., Jiang, W., and Inouye, M. (1998). Mol. Microbiol. 27, 873-874.

10. Zuker, M. and Stieger, P. (1982). *Nucleic Acids Res.* 9, 133-148
11. Inouye, M. (1983). Multipurpose expression cloning vehicles in *Escherichia coli*. In *Experimental Manipulation of Gene Expression*. Inouye, M. (ed.). New York, Academic Press, pp.15-32
- 5 12. Vind, J., Sorensen, M. A., Rasmussen, M. D., and Pedersen, S. (1993). *J. Mol. Biol.* 231, 678-688.
13. Iost, I and Dreyfus, M. (1995). *EMBO J.* 14, 3252-3261.
14. Wu, X-Q., Lyengar, P., and RajBhandary, U. L. (1996). *EMBO J.* 15, 4734-4739
15. Etchegaray, J.-P. and Inouye, M. (in press)
- 10 16. Winzeler, E., and Shapiro, L. (1997). *J. Bacteriol.* 179, 3981-3988
17. McCarthy, J. E. G., and Brimacombe, R. (1994). *TIG.* 10, 402-408
18. Faxen, M., Plumbridge, J. And Isaksson, L. A. (1991). *Nucleic Acids Res.* 19, 5247-5251
19. Nagai, N., Yuzawa, H., and Yura, T. (1991). *Proc. Natl. Acad. Sci. USA.* 88, 10515-10519
- 15 20. Ito, K., Kawakami, K., and Nakamura, Y. (1993). *Proc. Natl. Acad. Sci. USA.* 90, 302-306
21. Jiang, W., Fang, L., and Inouye, M. (1996).. *J. Bacteriol.* 178, 4919-4925
22. Harlocker, S. L., Rampersand, A., Yang, W.-P., and Inouye, M. (1993). *J. Bacteriol.* 175, 1956-1960
- 20 23. Miller, J. H. (1972). *Experiments in Molecular Genetics*. Cold Spring Harbor, NY: Cold Spring Harbor laboratory Press, pp. 352-355
24. Sarmientos, P., Sylvester, J. E., Contente, S., and Cashel, M. (1983). *Cell* 32, 1337-1346

25. Jones, P. G., Mitta, M., Kim, Y., Jiang, W., and Inouye, M. (1996). Proc. Natl. Acad. Sci. USA. 93, 76-80
26. Dammal, C. S., and Noller, H. F. (1995). Genes Dev. 9, 626-637
27. Ron, E. Z., Kohler, R. E., and Davis, B. D. (1966). Science 153, 1119-1120
- 5 28. Inouye, S., and Inouye, M. (1991). Directed Mutagenesis: A Practical Approach. M. J. McPherson, IRL Press, pp. 202
29. Altuvia, S., D. Kornitzer, D. Teff, and A. B. Oppenheim. 1989. Alternative mRNA structures of cIII gene of bacteriophage  $\lambda$  determine the rate of its translation initiation. J. Mol. Biol. 210:265-280.
- 10 30. Bae, W., P. G. Jones, and M. Inouye. 1997. CspA, the major cold-shock protein of *Escherichia coli*, negatively regulates its own gene expression. J. Bacteriol. 179:7081-7088.
31. Bae, W., S. Phadtare, K. Severinov, and M. Inouye. 1999. Characterization of *Escherichia coli* cspE, whose product negatively regulates transcription of cspA, the gene for the major cold shock protein. Mol. Microbiol. 31:1429-1441.
- 15 32. Brandi, A., P. Pietroni, C. O. Gualerzi, and C. L. Pon. 1996. Post-transcriptional regulation of CspA expression in *Escherichia coli*. Mol. Microbiol. 19:231-240.
33. Brosius, J., M. L. Palmer, P. J. Kennedy, and H. F. Noller. 1978. Complete nucleotide sequence of a 16S ribosomal RNA gene from *Escherichia coli*. Proc. Natl. Acad. Sci. USA 75:4801-4805.
- 20 34. Dontsova, O., S. Dokudovskaya, A. Kopylov, A. Bogdanov, J. Rinke-Appel, N. Junke, and R. Brimacombe. 1992. Three widely separated positions in the 16S rRNA lie in or close to the ribosomal decoding region: a site-directed cross-linking study with mRNA analogues. EMBO J. 11:3105-3116.
- 25 35. Etchegaray, J. P., and M. Inouye. 1999. CspA, CspB, and CspG, major cold shock proteins of *Escherichia coli*, are induced at low temperature under conditions that completely block protein synthesis. J. Bacteriol. 181:1827-1830.

36. Etchegaray, J. P., and M. Inouye. 1999. Translational enhancement by an element downstream of the initiation codon in *Escherichia coli*. J. Biol. Chem. 274: in press.
37. Etchegaray, J. P., P. G. Jones, and M. Inouye. 1996. Differential thermoregulation of two highly homologous cold-shock genes, *cspA* and *cspB*, of *Escherichia coli*. Genes Cells 1:171-178.
38. Fang, L., W. Jiang, W. Bae, and M. Inouye. 1997. Promoter-independent cold-shock induction of *cspA* and its derepression at 37°C by mRNA stabilization. Mol. Microbiol. 23:355-364.
39. Feng, W., R. Tejero, D. E. Zimmerman, M. Inouye, and G. T. Montelione. 1998. Solution NMR structure and backbone dynamics of the major cold-shock protein (CspA) from *Escherichia coli*: evidence for conformational dynamics in the single-stranded RNA-binding site. Biochemistry 37:10881-10896.
40. Goldenberg, D., I. Azar, and A. B. Oppenheim. 1996. Differential mRNA stability of the *cspA* gene in the cold-shock response of *Escherichia coli*. Mol. Microbiol. 19:241-248.
41. Goldenberg, D., I. Azar, and A. B. Oppenheim, A. Brandi, C. L. Pon, and C. O. Gualerzi. 1997. Role of *Escherichia coli cspA* promoter sequence and translational apparatus adaptation in the cold shock response. Mol. Gen. Genet. 256:282-290.
42. Goldstein, J., N. S. Pollitt, and M. Inouye. 1990. Major cold shock proteins of *Escherichia coli*. Proc. Natl. Acad. Sci. USA 87:283-287.
43. Hoe, N. P., and J. D. Goguen. 1993. Temperature sensing in *Yersinia pestis*: translation of the LcrF activator protein is thermally regulated. J. Bacteriol. 175:7901-7909.
44. Jiang, W., Y. Hou, and M. Inouye. 1997. CspA, the major cold-shock protein of *Escherichia coli*, is an RNA chaperone. J. Biol. Chem. 272:196-202.

45. Jones, P. G., and M. Inouye. 1996. RbfA, 30S ribosomal binding factor, is a cold-shock protein whose absence triggers the cold-shock response. *Mol. Microbiol.* 21:1207-1218.
46. Jones, P. G., R. A. VanBogelen, and F. C. Neidhardt. 1987. Induction of proteins in response to low temperature in *Escherichia coli*. *J. Bacteriol.* 169:2092-2095.
47. Lee, S. J., A. Xie, W. Jiang, J.-P. Etchegaray, P. G. Jones, and M. Inouye. 1994. Family of the major cold-shock protein, CspA (CS7.4) of *Escherichia coli*, whose members show a high sequence similarity with the eukaryotic Y-box binding proteins. *Mol. Microbiol.* 11:833-839.
48. Lopilato, J., S. Bortner, and J. Beckwith. 1986. Mutations in a new chromosomal gene of *Escherichia coli* K-12, *pcnB*, reduced plasmid copy number of pBR322 and its derivatives. *Mol. Gen. Genet.* 205:285-290.
49. Matsumoto, K., and A. P. Wolffe. 1998. Gene regulation by Y-box proteins: coupling control of transcription and translation. *Trends Cell Biol.* 8:318-323.
50. Miller, J. H. 1992. A Short Course in Bacterial Genetics - A Laboratory Manual and Handbook for *Escherichia coli* and Related Bacteria. Cold Spring Harbor Laboratory Press, Cold Harbor, NY.
51. Morita, M., M. Kanemori, H. Yanagi, and T. Yura. 1999. Heat-induced synthesis of  $\sigma^{32}$  in *Escherichia coli*: structural and functional dissection of *rpoH* mRNA secondary structure. *J. Bacteriol.* 181:401-410.
52. Morita, M. T., Y. Tanaka, T. S. Kodama, Y. Kyogoku, H. Yanagi, and T. Yura. 1999. Translational induction of heat shock transcription factor  $\sigma^{32}$ : evidence for a built-in RNA thermosensor. *Genes & Dev.* 13:655-665.
53. Nakashima, K., K. Kanamaru, T. Mizuno, and K. Horikoshi. 1996. A novel member of the *cspA* family of genes that is induced by cold-shock in *Escherichia coli*. *J. Bacteriol.* 178:2994-2997.



54. Newkirk, K., W. Feng, W. Jiang, R. Tejero, S. D. Emerson, M. Inouye, and G. T. Montelione. 1994. Solution NMR structure of the major cold shock protein (CspA) from *Escherichia coli*: identification of a binding epitope for DNA. *Proc. Natl. Acad. Sci. USA* 91:5114-5118.
55. Rinke-Appel, J., N. Junke, R. Brimacombe, S. Dokudovskaya, O. Dontsova, and A. Bogdanov. 1993. Site-directed cross-linking of mRNA analogues to 16S ribosomal RNA: a complete scan of cross-links from all positions between '+1' and '+16' on the mRNA, downstream from the decoding site. *Nucleic Acids Res.* 21:2853-2859.
56. Ross, W., K. K. Gosink, J. Salmon, K. Igarashi, C. Zou, A. Ishihama, K. Severinov, and R. L. Gourse. 1993. A third recognition element in bacterial promoters: DNA binding by the  $\alpha$  subunit of RNA polymerase. *Science* 262:1407-1413.
57. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning, a Laboratory Manual*, 2nd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
58. Sanger, F., S. Nickelen, and A. R. Coulson. 1977. DNA sequencing with chain-termination inhibitors. *Proc Natl Acad Sci USA* 74: 5463-5467.
59. Schindelin, H., W. Jiang, M. Inouye, and U. Heinemann. 1994. Crystal structure of CspA, the major cold shock protein of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 91:5119-5123.
60. Simons, R. W., F. Houman, and N. Kleckner. 1987. Improved single and multicopy lac-based cloning vectors for protein and operon fusions. *Gene* 53:85-96.
61. Storz, G. 1999. An RNA thermometer. *Genes & Dev.* 13:633-636.
62. Tanabe, H., J. Goldstein, M. Yang, and M. Inouye. 1992. Identification of the promoter region of the *Escherichia coli* major cold shock gene, *cspA*. 174:3867-3873.
63. Thieringer, H. A., P. G. Jones, and M. Inouye. 1998. Cold shock and adaptation. *BioEssay* 20:49-57.
64. Wang, N., K. Yamanaka, and M. Inouye. 1999. CspI, the ninth member of the CspA family of *Escherichia coli*, is induced upon cold shock. *J. Bacteriol.* 181:1603-1609.

65. Yamanaka, K., L. Fang, M. Inouye. 1998. The CspA family in *Escherichia coli*: multiple gene duplication for stress adaptation. *Mol. Microbiol.* 27:247-255.
66. Yamanaka, K., and M. Inouye. 1997. Growth-phase dependent expression of *cspD*, encoding a member of the CspA family in *Escherichia coli*. *J. Bacteriol.* 179:5126-5130.
67. Yuzawa, H., H. Nagai, H. Mori, and T. Yura. 1993. Heat induction of  $\sigma^{32}$  synthesis mediated by mRNA secondary structure: a primary step of the heat shock response in *Escherichia coli*. *Nucleic Acids Res.* 21:5449-5455.
68. Doniger, J., D.Landsman, M.A.Gonda, and G.Wistow. 1992. The product of *unr*, the highly conserved gene upstream of *N-ras*, contains multiple repeats similar to the cold-shock domain (CSD), a putative DNA-binding motif. *The New Biologist* 4:389-396.
69. Etchegaray, J.P., P.G.Jones, and M.Inouye. 1996. Differential thermoregulation of two highly homologous cold-shock genes, *cspA* and *cspB*, of *Escherichia coli*. *Genes to Cells.* in press.
70. Inouye, S., X. Soheron, T. Franceschini, K. Nakamura, K. Hakura, and M. Inouye. 1982. Role of positive charge on the amino-terminal region of the signal peptide in protein secretion across the membrane. *Proc. Natl. Acad. Sci. USA* 79:3438-3441.
71. Jiang, W., P.G. Jones, and M. Inouye. 1993. Chloramphenicol induces the transcription of the major cold-shock gene of *Escherichia coli*, *cspA*. *J. Bacteriol.* 175:5824-5828.
72. Jones, P.G., M. Cashel, G.Glaser, and F.C. Neidhardt. 1992. Function of a relaxed-like state following temperature downshifts in *Escherichia coli*. *J. Bacteriol.* 174:3903-3904.
73. Jones, P.G. and M. Inouye. 1994. The cold-shock response-a hot topic. *Mol. Microbiol.* 11:811-818.

74. Jones, P.G., M. Mitta, Y. Kim, W. Jiang, and M. Inouye. 1995. Cold shock induces a new major ribosomal-associated protein which unwinds double-stranded RNA in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 93:76-80.
75. Lerner, C.G. and M. Inouye. 1990. Low copy number plasmids for regulated low-level expression of cloned genes in *Escherichia coli* with blue/white insert screening capability. *Nucl. Acids Res.* 18:4631.
76. Toone, W.M., K.E. Rudd, and J.D. Friesen. 1991. dead, a new *Escherichia coli* gene encoding a presumed ATP-dependent RNA helicase, can suppress a mutation in *rpsB*, the gene encoding ribosomal protein S2. *J. Bacteriol.* 173:3291-3302.
77. Vieira, J. and J. Messing. 1982. The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers. *Gene* 19:259-268.
78. Wolffe, A.P., S. Tafuri, M. Ranjan, and M. Familiar. 1992. The Y-box factors: a family of nucleic acid binding protein conserved from *Escherichia coli* to man. *New Biol.* 4:290-298.
79. Wolffe, A.P. 1994. Structural and functional properties of the evolutionarily ancient Y-box family of nucleic acid binding proteins. *BioEssays.* 16:245-251.
80. Yamanaka, K., T. Mitani, T. Ogura, H. Niki, and S. Hiraga. 1994. Cloning, sequencing, and characterization of multicopy suppressors of a *mukB* mutation in *Escherichia coli*. *Mol. Microbiol.* 13:301-312.

We claim:

1. An isolated nucleic acid molecule that prolongs the expression of cold shock inducible genes under conditions that elicit the cold shock response in a bacterium.
2. The nucleic acid molecule of Claim 1, wherein said molecule comprises a 5'-UTR  
5 of a cold shock inducible gene or a substantially homologous sequence thereof.
3. The nucleic acid molecule of Claim 2, wherein said 5'-UTR is a 5'-UTR of a cold-shock inducible gene selected from the group consisting of *cspA*, *cspB* and *csdA*.
4. The isolated nucleic acid molecule of Claim 2, wherein said 5'-UTR comprises a cold box or a substantially homologous sequence thereof.
- 10 5. The nucleic acid molecule of Claim 3, wherein said 5'-UTR comprises nucleotides +1 to +11 of the *cspA* 5'-UTR or a substantially homologous sequence thereof.
6. The nucleic acid molecule of Claim 1, wherein said cold shock inducible gene interacts with CspA protein.
7. An isolated nucleic acid molecule that represses the expression of cold shock inducible genes under physiological conditions.
- 15 8. The isolated nucleic acid molecule of Claim 7, comprising at least a portion of the 5'-UTR of a cold shock inducible gene.
9. The isolated nucleic acid molecule of Claim 8, wherein said cold-shock inducible gene is selected from the group consisting of *cspA*, *cspB*, and *csdA*.
- 20 10. The nucleic acid molecule of Claim 8 wherein said cold-shock inducible gene comprises nucleotides +56 to +117 of *cspA* or nucleotides having substantial homology to nucleotides +56 to +117 of *cspA*.
11. A non-coding nucleic acid molecule that enhances the translation of cold shock inducible genes under conditions that elicit the cold shock response of a bacterium.

12. The nucleic acid molecule of Claim 11 comprising at least a portion of the 5'-UTR of a cold shock inducible gene.

13. The nucleic acid molecule of Claim 12 wherein said cold shock inducible gene is selected from the group consisting of *cspA*, *cspB*, and *csdA*.

14. The nucleic acid molecule of Claim 13, comprising nucleotides +123 to +135 of *cspA* or nucleotides having substantial homology to +123 to +135 of *cspA*.

15. The nucleic acid molecule of Claim 14 comprising having a sequence selected from the group consisting of SEQ ID NO:48, SEQ ID NO:49, and SEQ ID NO:50.

16. A nucleic acid vector comprising a downstream box and a nucleic acid molecule that enhances translation of cold shock inducible genes under conditions that elicit the cold shock response in a bacterium.

17. The construct of Claim 16 further comprising a Shine-Dalgarno sequence.

18. A nucleic acid vector wherein at least a portion of said vector comprises a cold box, a translational enhancer, and a downstream box; said vector directing prolonged expression and enhancing translation of a gene under conditions that elicit a cold shock response in a bacterium.

19. A nucleic acid vector that directs the prolonged expression and enhances the translation of a gene under conditions of physiological stress that elicit the cold shock response of a bacterium, and represses the expression of the gene under physiological conditions comprising a cold box, at least a portion of the 5'-UTR of a cold-shock inducible gene that represses the expression of cold-shock inducible genes, at least a portion of the 5'-UTR of a cold-shock inducible gene that enhances translation of cold-shock genes, and a downstream box.

20. The vector of claim 16 wherein said the gene is a cold shock inducible gene.

21. The vector of Claim 18, wherein said gene is a cold-shock inducible gene.

22. The vector of Claim 19, wherein said gene is a cold-shock inducible gene.
23. The vector of Claim 16 wherein said gene is a heterologous gene.
24. The vector of Claim 18 wherein said gene is a heterologous gene.
25. The vector of Claim 19, wherein said gene is a heterologous gene.
- 5 26. The vector of Claim 16, further comprising a promoter and at least one restriction site downstream of said 5' UTR and said downstream box for inserting an additional DNA fragment.
27. The vector of Claim 18, further comprising a promoter and at least one restriction site downstream of said 5' UTR and said downstream box for inserting an additional DNA fragment.
- 10 28. The vector of Claim 19, further comprising a promoter and at least one restriction site downstream of said 5' UTR and said downstream box for inserting an additional DNA fragment.
29. The vector of claim 26, wherein said additional DNA fragment encodes a cold shock inducible gene.
- 15 30. The vector of 27, wherein said additional DNA fragment encodes a cold shock inducible gene.
31. The vector of Claim 28, wherein said additional DNA fragment encodes a cold shock inducible gene.
- 20 32. The vector of Claim 26, wherein said additional DNA fragment encodes a heterologous gene.
33. The vector of Claim 27, wherein said additional DNA fragment encodes a heterologous gene.
- 25 34. The vector of Claim 28, wherein said additional DNA fragment encodes a heterologous gene.

35. A transformed bacteria containing the vector of Claim 16.
36. A transformed bacteria containing the vector of Claim 18.
37. A transformed bacteria containing the vector of Claim 19.
38. A method for overexpressing a gene comprising transforming bacteria with a  
5 nucleic acid vector of Claim 16 and subjecting said bacteria to conditions that elicit a  
cold shock response.
39. A method for overexpressing a gene comprising transforming bacteria with a  
nucleic acid vector of Claim 18 and subjecting said bacteria to conditions that elicit a  
cold shock response.
40. A method for overexpressing a gene comprising transforming bacteria with a  
10 nucleic acid vector of Claim 19 and subjecting said bacteria to conditions that elicit a  
cold shock response.
41. The method of Claim 37, wherein said overexpression causes a reduction in the  
expression of at least one endogenous protein.
42. The method of Claim 38, wherein said overexpression causes a reduction in the  
15 expression of at least one endogenous protein.
43. The method of Claim 39, wherein said overexpression causes a reduction in the  
expression of at least one endogenous protein.
44. The method of Claim 37, wherein said conditions that elicit a cold shock response  
20 comprise subjecting said bacteria to a sufficiently low temperature to elicit a cold-  
shock response.
45. The method of Claim 38, wherein said conditions that elicit a cold shock response  
comprise subjecting said bacteria to a sufficiently low temperature to elicit a cold-  
shock response.

46. The method of Claim 39, wherein said conditions that elicit a cold shock response comprise subjecting said bacteria to a sufficiently low temperature to elicit a cold-shock response.
47. The method of Claim 43, wherein said temperature is about 10-15°C.
48. The method of Claim 44, wherein said temperature is about 10-15°C.
49. The method of Claim 45, wherein said temperature is about 10-15°C.
50. A vector capable of expressing a heterologous gene in a bacterium at physiological temperature or under conditions that elicit a cold shock response comprising in the following order: a promoter, at least a portion of a 5'-UTR of a cold shock protein gene, a Shine-Dalgarno sequence, a translational initiation codon, a downstream box of a cold shock inducible gene, and at least one restriction enzyme recognition site for insertion of said heterologous gene.
51. The vector of Claim 50, further comprising an additional nucleic acid fragment wherein said fragment encodes a cold shock inducible gene.
52. The vector of Claim 50, further comprising an additional nucleic acid wherein said fragment encodes a heterologous gene.
53. A transformed bacteria containing the vector of Claim 50.
54. A method of overexpressing a gene comprising transforming bacteria with a nucleic acid vector of Claim 50 and subjecting said bacteria to conditions that elicit a cold shock response.
55. The method of Claim 54, wherein said conditions that elicit a cold shock response comprise subjecting said bacteria to a sufficiently low temperature to elicit a cold-shock response.
56. The method of Claim 55, wherein said temperature is about 10-15°C.






FIG. 1A




FIG. 1B




FIG. 2




FIG. 3

100  
 80  
 60  
 40  
 20  
 0




FIG. 4






FIG. 6E



FIG. 6C




FIG. 6A




FIG. 6A




FIG. 6B




FIG. 6C






FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 11A

FIG. IIB





FIG. IIC








FIG. 12




FIG. 13A




FIG. 13B




FIG. 14A  
pMM67






FIG. 14C  
pMM023




FIG. 14D  
pMM024




FIG. 14E  
pMM025






FIG. 14F  
pMM026




FIG. 15A




FIG. 15B




FIG. 15C




FIG. 15D




FIG. 15E




FIG. 16A




FIG. 16B





FIG. 16C




FIG. 16D




FIG. 17A




FIG. 17B




FIG. 17C




FIG. 17D




FIG. 17E




FIG. 17F




FIG. 17G




FIG. 17H






FIG. 18A

---




FIG. 18B




FIG. 19A




FIG. 19B




FIG. 19C




FIG. 20A




FIG. 20B

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

**DECLARATION FOR UTILITY OR  
DESIGN  
PATENT APPLICATION  
(37 CFR 1.63)**

☐ Declaration Submitted with Initial Filing

OR

☒ Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)

Attorney Docket Number 913.6600CIP

First Named Inventor Li Fang

**COMPLETE IF KNOWN**

Application Number

Filing Date

April 19, 2000

Group Art Unit

Examiner Name

As a below named inventor, I hereby declare

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**COLD-SHOCK REGULATORY ELEMENTS, CONSTRUCTS THEREOF, AND METHODS OF USE**

the specification of which

(Title of the invention)

☐ is attached hereto

OR

☒ was filed on (MM/DD/YYYY)

April 19, 2000

as United States Application Number or PCT International

Application Number

and was amended on (MM/DD/YYYY)

(if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached? YES NO
			<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.
60/096,938 60/143,380	August 20, 1998	

[Page 1 of 2]

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)  
Approved for use through 9/30/00, OMB 0651-0032  
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE  
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

## DECLARATION—Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.55 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U. S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)
PCT/US99/19030 09/293,427 08/769,945 08/203,806	April 20, 1999 April 16, 1999 December 19, 1996 March 1, 1994	

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

☒ Customer Number

022469

OR

☐ Registered practitioner(s) name/registration number listed below

Place Customer Number Bar Code Label here

Name	Registration Number	Name	Registration Number

☐ Additional registered practitioner(s) named on supplemental Registered Practitioner information sheet PTO/SB/02C attached hereto.

Direct all correspondence to:

☒ Customer Number or

022469

OR

☐ Correspondence address below

Name	Gerard Weiser - IP Department				
Address	Schnader Harrison Segal & Lewis LLP				
Address	1600 Market Street, Suite 3600				
City	Philadelphia	State	PA	ZIP	19103
Country	US	Telephone	215-751-2427	Fax	215-751-2658

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))		Family Name or Surname	
Li	Fang		
Inventor's Signature		Date	
Residence: City	New York	State	NY
		Country	US
Post Office Address	Apartment 883, 50 98th Street		
Post Office Address			
City	New York	State	NY
		ZIP	10029
		Country	US

☒ Additional inventors are being named on the 1 supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto





Please type a plus sign (+) inside this box → +

PTO/SB/02A (3-97)

Approved for use through 9/30/98. OMB 0551-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

+

<b>DECLARATION</b>	<b>ADDITIONAL INVENTOR(S) Supplemental Sheet</b> Page <u>1</u> of <u>2</u>
--------------------	---

<b>Name of Additional Joint Inventor, if any:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle (if any))				Family Name or Surname			
Weining				Jiang			
Inventor's Signature		Date					
Residence: City		New York		State		NY	
				Country		US	
Post Office Address		Apartment 2H, 430 East 63rd Street					
Post Office Address							
City		New York		State		NY	
				ZIP		10021	
				Country		US	
<b>Name of Additional Joint Inventor, if any:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle (if any))				Family Name or Surname			
Masanori				Mitta			
Inventor's Signature		Date					
Residence: City		Kyoto		State		Japan	
				Country		Japan	
Post Office Address		42-14-2-321, Hensyoware, Jusauchi, Tanabe-cho, Tsuzuki-gun					
Post Office Address							
City		Kyoto		State		Japan	
				ZIP		610-03	
				Country		Japan	
<b>Name of Additional Joint Inventor, if any:</b>		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle (if any))				Family Name or Surname			
Masayori				Inouye			
Inventor's Signature		Date					
Residence: City		Piscataway		State		NJ	
				Country		US	
Post Office Address		Apartment 107B, 280 Riber Road					
Post Office Address							
City		Piscataway		State		NJ	
				ZIP		08854	
				Country		US	

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

+

Please type a plus sign (+) inside this box → ☐

PTO/SB/02A (3-97)

Approved for use through 9/30/98. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.



<b>DECLARATION</b>	<b>ADDITIONAL INVENTOR(S)</b> <b>Supplemental Sheet</b> Page <u>2</u> of <u>2</u>
--------------------	---

<b>Name of Additional Joint Inventor, if any:</b>	<input type="checkbox"/> A petition has been filed for this unsigned inventor
---	---

Given Name (first and middle [if any])	Family Name or Surname
Jean-Pierre	Etchegaray

Inventor's Signature		Date	
----------------------	--	------	--

Residence: City	Highland Park	State	NY	Country	US	Citizenship
-----------------	---------------	-------	----	---------	----	-------------

Post Office Address	223 Lincoln Avenue
---------------------	--------------------

Post Office Address	
---------------------	--

City	Highland Park	State	NJ	ZIP	08904	Country	US
------	---------------	-------	----	-----	-------	---------	----

<b>Name of Additional Joint Inventor, if any:</b>	<input type="checkbox"/> A petition has been filed for this unsigned inventor
---	---

Given Name (first and middle [if any])	Family Name or Surname

Inventor's Signature		Date	
----------------------	--	------	--

Residence: City		State		Country		Citizenship
-----------------	--	-------	--	---------	--	-------------

Post Office Address	
---------------------	--

Post Office Address	
---------------------	--

City		State		ZIP		Country	
------	--	-------	--	-----	--	---------	--

<b>Name of Additional Joint Inventor, if any:</b>	<input type="checkbox"/> A petition has been filed for this unsigned inventor
---	---

Given Name (first and middle [if any])	Family Name or Surname

Inventor's Signature		Date	
----------------------	--	------	--

Residence: City		State		Country		Citizenship
-----------------	--	-------	--	---------	--	-------------

Post Office Address	
---------------------	--

Post Office Address	
---------------------	--

City		State		ZIP		Country	
------	--	-------	--	-----	--	---------	--

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.



IN THE EUROPEAN PATENT OFFICE  
BEFORE THE INTERNATIONAL SEARCHING AUTHORITY

Atty. Docket No: 913.6600PCT

In re International patent application of

THE UNIVERSITY OF MEDICINE AND DENTISTRY et al.

International Application No. PCT/US99/19030

International Filing Date: August 20, 1999

For: COLD-SHOCK REGULATORY ELEMENTS, CONSTRUCTS THEREOF, AND  
METHODS OF USE

European Patent Office  
Storage and Retrieval of Amino  
Acid and Nucleotide Data  
Room POH09  
Patentlaan 2  
P.B. 5818  
NL-2280 HV Rijswijk  
The Netherlands


STATEMENT ACCOMPANYING SEQUENCE LISTING

Dear Sir:

The undersigned hereby states that the Sequence Listing submitted concurrently herewith does not include matter which goes beyond the content of the application as filed and that the information recorded on the diskette submitted concurrently herewith is identical to the written Sequence Listing.

Respectfully submitted,

Dec. 17, 1999  
Date

  
James A. Coburn

HARBOR CONSULTING  
Intellectual Property Services  
1500A Lafayette Road  
Suite 262  
Portsmouth, N.H.  
(800) 318-3021

## SEQUENCE LISTING

<110> THE UNIVERSITY OF MEDICINE AND DENTISTRY

<120> COLD-SHOCK REGULATORY ELEMENTS, CONSTRUCTS THEREOF, AND METHODS OF USE

<130> 913.6600PCT

<140> PCT/US99/19030

<141> 1999-08-20

<160> 71

<170> PatentIn Ver. 2.0

<210> 1  
<211> 14  
<212> RNA  
<213> E. coli

<400> 1  
acuuugugau ucau 14

<210> 2  
<211> 14  
<212> RNA  
<213> E. coli

<400> 2  
augacuggua ucgu 14

<210> 3  
<211> 14  
<212> RNA  
<213> E. coli

<400> 3  
augacugguu ucgu 14

<210> 4  
<211> 14  
<212> RNA  
<213> E. coli

<400> 4  
augacugguu uagu 14

<210> 5  
<211> 14  
<212> RNA  
<213> E. coli

<400> 5  
augaguuauug uaga 14

<210> 6  
<211> 14

<212> RNA  
<213> E. coli

<400> 6  
auggcgaaaa gaau

14

<210> 7  
<211> 47  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: mRNA construct

<220>  
<223> n = g, c, u or a

<220>  
<223> This sequence may encompass a construct wherein  
the "n" region may be 0-30.

<400> 7  
augnnnnnnnn nnnnnnnnnn nnnnaugacug guaucgu

47

<210> 8  
<211> 47  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: DNA which  
encodes for the mRNA construct

<220>  
<223> n = g, c, t, or a

<220>  
<223> This sequence may encompass a construct wherein  
the "n" region may be 0-30.

<400> 8  
atgnnnnnnnn nnnnnnnnnn nnnnatgactg gtagcgt

47

<210> 9  
<211> 15  
<212> DNA  
<213> E. coli

<220>  
<221> MOD\_RES  
<220> (5)  
<223> a substituted by t

<220>  
<221> MOD\_RES  
<220> (6)  
<223> t substituted by c

```

<220>
<221> MOD_RBS
<220> (9)
<223> a substituted by g

<400> 9
aattmntana ggtaa                                15

<210> 10
<211> 15
<212> DNA
<213> E. coli

<400> 10
acggttctag acgta                                15

<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 11
cggcattaag taagcagttg                            20

<210> 12
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 12
ctggatcctt taatggtctg tacgtcaaac cgt            33

<210> 13
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 13
cggaattcag cctgtaatct ct                          22

<210> 14
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 14

```

ctgtcgactt acttacggcg ttgc 24

<210> 15  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 15  
 gacaggatta aaaatcgag 19

<210> 16  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 16  
 aaccgttgat gtgca 15

<210> 17  
 <211> 28  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 17  
 ccttgctagc cgattaatca taaatatg 28

<210> 18  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 18  
 ccggtaccag gttgaacat ttt 23

<210> 19  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 19  
 actacacttt gatgtgcatt agc 23

<210> 20

<211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 20  
 caacgataag cttaaatggt ctgt

24

<210> 21  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 21  
 taaaggctct tgaagggact t

21

<210> 22  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 22  
 cggcgatata atgtgcacta cgaggg

26

<210> 23  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 23  
 tacctttaag gogtgcttta cagatt

26

<210> 24  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 24  
 gcacatcaaa gtgtagtaag gcaa

24

<210> 25  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence



```

<220>
<223> Description of Artificial Sequence: Primer

<400> 25
taaagcttat cgttgatacc c                               21

<210> 26
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 26
tcaagagcct ttaacgcttc aaaa                             24

<210> 27
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 27
gcacattata tcgccgaaag gc                               22

<210> 28
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 28
aaagcacgcc ttaaaggtaa tacact                           26

<210> 29
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
      Oligonucleotide #8509

<400> 29
ctagccgaaa ggcacaaatt aagaggtat taataatgaa agggggaatt cca       53

<210> 30
<211> 53
<212> DNA
<213> Artificial Sequence

<220>

```

<223> Description of Artificial Sequence:  
Oligonucleotide #8510

<400> 30  
agcttggaat tccccctttc attattaata ccctcttaat ttgtgccttt cgg 53

<210> 31  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 31  
cggatccag cttaataata gct 23

<210> 32  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 32  
cggatccag attgacatt ctaca 25

<210> 33  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 33  
ccgatccag gttaaaccat ttt 23

<210> 34  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 34  
ccgatccag acctttatca gcgtt 25

<210> 35  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 35  
 gaaaggtctca agttacttca tgtagaatg 29

<210> 36  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 36  
 cattctacat gaagtaactt gaggcctttc 29

<210> 37  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide

<400> 37  
 aattaatcac aaagtggg 18

<210> 38  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide

<400> 38  
 aattcccact ttgtgatt 18

<210> 39  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide

<400> 39  
 aattatgaat cacaaagtgg g 21

<210> 40  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide

<400> 40  
 aattcccaact ttgtgattca t 21  
  
 <210> 41  
 <211> 48  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide  
  
 <400> 41  
 ctagccctta ttaataatga aagggggaat tatgaatcac aaagtggg 48  
  
 <210> 42  
 <211> 48  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide  
  
 <400> 42  
 aattcccaact ttgtgattca taattccccc ttctattatt aataaggg 48  
  
 <210> 43  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide  
  
 <400> 43  
 ctagccctta ttaataatga atcacaaagt ggg 33  
  
 <210> 44  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: annealed  
 oligonucleotide  
  
 <400> 44  
 aattcccaact ttgtgattca ttattaataa ggg 33  
  
 <210> 45  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>

<223> Description of Artificial Sequence: annealed  
oligonucleotide

<400> 45  
ctagagggtta ttaataatga atcacaaagt ggg 33

<210> 46  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: annealed  
oligonucleotide

<400> 46  
aattccact ttgtgattca ttattaatac cct 33

<210> 47  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 47  
cgccagggtt ttccagtcga cgac 24

<210> 48  
<211> 13  
<212> RNA  
<213> E. coli

<400> 48  
gccgaaaggc aca 13

<210> 49  
<211> 13  
<212> RNA  
<213> E. coli

<400> 49  
gccgaaaggc uca 13

<210> 50  
<211> 13  
<212> RNA  
<213> E. coli

<400> 50  
gccgaaaggc cca 13

<210> 51  
<211> 47  
<212> RNA  
<213> E. coli

<400> 51  
 uugacaacca cggaaguuuu cagagaugag aaugugccuu cgggaaac 47  
  
 <210> 52  
 <211> 40  
 <212> RNA  
 <213> E. coli  
  
 <400> 52  
 gccgaaaggc acacuuuuuu auuaaaggua auacacuaug 40  
  
 <210> 53  
 <211> 32  
 <212> RNA  
 <213> E. coli  
  
 <400> 53  
 gccgaaaggc ucaaguuuag gaauugagaa ug 32  
  
 <210> 54  
 <211> 34  
 <212> RNA  
 <213> E. coli  
  
 <400> 54  
 gccgaaaggc ccaaaaugaa ggaaguaaaa uaug 34  
  
 <210> 55  
 <211> 162  
 <212> RNA  
 <213> E. coli  
  
 <400> 55  
 acgguuugac guacagacca uuaaagcagu guaguaaggc aagucccuuc aagaguuauc 60  
 guugauaccc cucguagugc acauuccuuu aacgcuucaa aaucuguaaa gcacgccaaua 120  
 ucgcgcaag gcacacuuua uuauuaaagg uauuacacua ug 162  
  
 <210> 56  
 <211> 136  
 <212> RNA  
 <213> E. coli  
  
 <400> 56  
 aaguguaqua aggcagucc cuucaagagu uaucguugau accccucgua gucacauuc 60  
 cuuuacgcgu ucaaaucug uaaagcacgc cauaucgccg aaaggcacac uuaauuuuaa 120  
 aagguaauac acuauug 136  
  
 <210> 57  
 <211> 134  
 <212> RNA  
 <213> E. coli  
  
 <400> 57  
 acgguuugac guacagacca uuaaagcuua ucguugauac cccucguagu gcacauuccu 60  
 uuaacgcuuc aaaaucugua aagcacgcca uaucgccgaa aggcacacuu aauuuuuuaa 120  
 gguauuacac uaug 134  
  
 <210> 58

<211> 131  
 <212> RNA  
 <213> E. coli

<400> 58  
 acgguuugac guacagacca uuaaagcagu guaguaaggc aagucccuuc aagagcuua 60  
 acgcuucaa aucuguaag cagcccauau cgccgaaagg cacacuuaa uauuaagggu 120  
 aauacacuau g 131

<210> 59  
 <211> 130  
 <212> RNA  
 <213> E. coli

<400> 59  
 acgguuugac guacagacca uuaaagcagu guaguaaggc aagucccuuc aagaguuuac 60  
 guugauaccc cucguagugc acuuuuuau gccgaaaggc acacuuaauu auuaaaggua 120  
 auacacuau g 130

<210> 60  
 <211> 136  
 <212> RNA  
 <213> E. coli

<400> 60  
 acgguuugac guacagacca uuaaagcagu guaguaaggc aagucccuuc aagaguuuac 60  
 guugauaccc cucguagugc acuuuccuuu aacgcuucaa aaucuguaaa gcacgcuua 120  
 aagguauuac acuaug 136

<210> 61  
 <211> 65  
 <212> RNA  
 <213> E. coli

<400> 61  
 uuaaggaug uagauguca auaaaauga cugguuuagu aaaaugguuu aacgcugaua 60  
 aaggu 65

<210> 62  
 <211> 39  
 <212> RNA  
 <213> E. coli

<400> 62  
 cuuaaccuuc gggagggcgc uuaccacuuu gugauuau 39

<210> 63  
 <211> 15  
 <212> RNA  
 <213> E. coli

<400> 63  
 uacuuagugu uucac 15

<210> 64  
 <211> 12  
 <212> RNA  
 <213> E. coli

<400> 64  
 aaucacaaag ug 12  
  
 <210> 65  
 <211> 15  
 <212> RNA  
 <213> E. coli  
  
 <400> 65  
 augaauca aagug 15  
  
 <210> 66  
 <211> 47  
 <212> RNA  
 <213> E. coli  
  
 <400> 66  
 ucuagaggu auuaauaag aaagggggaa uuccaagcuu ggaucg 47  
  
 <210> 67  
 <211> 15  
 <212> RNA  
 <213> E. coli  
  
 <400> 67  
 cacuuuguga uucau 15  
  
 <210> 68  
 <211> 48  
 <212> RNA  
 <213> E. coli  
  
 <400> 68  
 ucuagaggu auuaauaag aaagggggaa uuaugaauc caaagugg 48  
  
 <210> 69  
 <211> 48  
 <212> RNA  
 <213> E. coli  
  
 <400> 69  
 ucuagcccu auuaauaag aaagggggaa uuaugaauc caaagugg 48  
  
 <210> 70  
 <211> 33  
 <212> RNA  
 <213> E. coli  
  
 <400> 70  
 ucuagcccu auuaauaag aaucacaaag ugg 33  
  
 <210> 71  
 <211> 33  
 <212> RNA  
 <213> E. coli



&lt;400&gt; 71

ucuagaggggu auuaauaaug aaucacaaag ugg

33

1100  
 1101  
 1102  
 1103  
 1104  
 1105  
 1106  
 1107  
 1108  
 1109  
 1110  
 1111  
 1112  
 1113  
 1114  
 1115  
 1116  
 1117  
 1118  
 1119  
 1120  
 1121  
 1122  
 1123  
 1124  
 1125  
 1126  
 1127  
 1128  
 1129  
 1130  
 1131  
 1132  
 1133  
 1134  
 1135  
 1136  
 1137  
 1138  
 1139  
 1140  
 1141  
 1142  
 1143  
 1144  
 1145  
 1146  
 1147  
 1148  
 1149  
 1150  
 1151  
 1152  
 1153  
 1154  
 1155  
 1156  
 1157  
 1158  
 1159  
 1160  
 1161  
 1162  
 1163  
 1164  
 1165  
 1166  
 1167  
 1168  
 1169  
 1170  
 1171  
 1172  
 1173  
 1174  
 1175  
 1176  
 1177  
 1178  
 1179  
 1180  
 1181  
 1182  
 1183  
 1184  
 1185  
 1186  
 1187  
 1188  
 1189  
 1190  
 1191  
 1192  
 1193  
 1194  
 1195  
 1196  
 1197  
 1198  
 1199  
 1200  
 1201  
 1202  
 1203  
 1204  
 1205  
 1206  
 1207  
 1208  
 1209  
 1210  
 1211  
 1212  
 1213  
 1214  
 1215  
 1216  
 1217  
 1218  
 1219  
 1220  
 1221  
 1222  
 1223  
 1224  
 1225  
 1226  
 1227  
 1228  
 1229  
 1230  
 1231  
 1232  
 1233  
 1234  
 1235  
 1236  
 1237  
 1238  
 1239  
 1240  
 1241  
 1242  
 1243  
 1244  
 1245  
 1246  
 1247  
 1248  
 1249  
 1250  
 1251  
 1252  
 1253  
 1254  
 1255  
 1256  
 1257  
 1258  
 1259  
 1260  
 1261  
 1262  
 1263  
 1264  
 1265  
 1266  
 1267  
 1268  
 1269  
 1270  
 1271  
 1272  
 1273  
 1274  
 1275  
 1276  
 1277  
 1278  
 1279  
 1280  
 1281  
 1282  
 1283  
 1284  
 1285  
 1286  
 1287  
 1288  
 1289  
 1290  
 1291  
 1292  
 1293  
 1294  
 1295  
 1296  
 1297  
 1298  
 1299  
 1300  
 1301  
 1302  
 1303  
 1304  
 1305  
 1306  
 1307  
 1308  
 1309  
 1310  
 1311  
 1312  
 1313  
 1314  
 1315  
 1316  
 1317  
 1318  
 1319  
 1320  
 1321  
 1322  
 1323  
 1324  
 1325  
 1326  
 1327  
 1328  
 1329  
 1330  
 1331  
 1332  
 1333  
 1334  
 1335  
 1336  
 1337  
 1338  
 1339  
 1340  
 1341  
 1342  
 1343  
 1344  
 1345  
 1346  
 1347  
 1348  
 1349  
 1350  
 1351  
 1352  
 1353  
 1354  
 1355  
 1356  
 1357  
 1358  
 1359  
 1360  
 1361  
 1362  
 1363  
 1364  
 1365  
 1366  
 1367  
 1368  
 1369  
 1370  
 1371  
 1372  
 1373  
 1374  
 1375  
 1376  
 1377  
 1378  
 1379  
 1380  
 1381  
 1382  
 1383  
 1384  
 1385  
 1386  
 1387  
 1388  
 1389  
 1390  
 1391  
 1392  
 1393  
 1394  
 1395  
 1396  
 1397  
 1398  
 1399  
 1400  
 1401  
 1402  
 1403  
 1404  
 1405  
 1406  
 1407  
 1408  
 1409  
 1410  
 1411  
 1412  
 1413  
 1414  
 1415  
 1416  
 1417  
 1418  
 1419  
 1420  
 1421  
 1422  
 1423  
 1424  
 1425  
 1426  
 1427  
 1428  
 1429  
 1430  
 1431  
 1432  
 1433  
 1434  
 1435  
 1436  
 1437  
 1438  
 1439  
 1440  
 1441  
 1442  
 1443  
 1444  
 1445  
 1446  
 1447  
 1448  
 1449  
 1450  
 1451  
 1452  
 1453  
 1454  
 1455  
 1456  
 1457  
 1458  
 1459  
 1460  
 1461  
 1462  
 1463  
 1464  
 1465  
 1466  
 1467  
 1468  
 1469  
 1470  
 1471  
 1472  
 1473  
 1474  
 1475  
 1476  
 1477  
 1478  
 1479  
 1480  
 1481  
 1482  
 1483  
 1484  
 1485  
 1486  
 1487  
 1488  
 1489  
 1490  
 1491  
 1492  
 1493  
 1494  
 1495  
 1496  
 1497  
 1498  
 1499  
 1500  
 1501  
 1502  
 1503  
 1504  
 1505  
 1506  
 1507  
 1508  
 1509  
 1510  
 1511  
 1512  
 1513  
 1514  
 1515  
 1516  
 1517  
 1518  
 1519  
 1520  
 1521  
 1522  
 1523  
 1524  
 1525  
 1526  
 1527  
 1528  
 1529  
 1530  
 1531  
 1532  
 1533  
 1534  
 1535  
 1536  
 1537  
 1538  
 1539  
 1540  
 1541  
 1542  
 1543  
 1544  
 1545  
 1546  
 1547  
 1548  
 1549  
 1550  
 1551  
 1552  
 1553  
 1554  
 1555  
 1556  
 1557  
 1558  
 1559  
 1560  
 1561  
 1562  
 1563  
 1564  
 1565  
 1566  
 1567  
 1568  
 1569  
 1570  
 1571  
 1572  
 1573  
 1574  
 1575  
 1576  
 1577  
 1578  
 1579  
 1580  
 1581  
 1582  
 1583  
 1584  
 1585  
 1586  
 1587  
 1588  
 1589  
 1590  
 1591  
 1592  
 1593  
 1594  
 1595  
 1596  
 1597  
 1598  
 1599  
 1600  
 1601  
 1602  
 1603  
 1604  
 1605  
 1606  
 1607  
 1608  
 1609  
 1610  
 1611  
 1612  
 1613  
 1614  
 1615  
 1616  
 1617  
 1618  
 1619  
 1620  
 1621  
 1622  
 1623  
 1624  
 1625  
 1626  
 1627  
 1628  
 1629  
 1630  
 1631  
 1632  
 1633  
 1634  
 1635  
 1636  
 1637  
 1638  
 1639  
 1640  
 1641  
 1642  
 1643  
 1644  
 1645  
 1646  
 1647  
 1648  
 1649  
 1650  
 1651  
 1652  
 1653  
 1654  
 1655  
 1656  
 1657  
 1658  
 1659  
 1660  
 1661  
 1662  
 1663  
 1664  
 1665  
 1666  
 1667  
 1668  
 1669  
 1670  
 1671  
 1672  
 1673  
 1674  
 1675  
 1676  
 1677  
 1678  
 1679  
 1680  
 1681  
 1682  
 1683  
 1684  
 1685  
 1686  
 1687  
 1688  
 1689  
 1690  
 1691  
 1692  
 1693  
 1694  
 1695  
 1696  
 1697  
 1698  
 1699  
 1700  
 1701  
 1702  
 1703  
 1704  
 1705  
 1706  
 1707  
 1708  
 1709  
 1710  
 1711  
 1712  
 1713  
 1714  
 1715  
 1716  
 1717  
 1718  
 1719  
 1720  
 1721  
 1722  
 1723  
 1724  
 1725  
 1726  
 1727  
 1728  
 1729  
 1730  
 1731  
 1732  
 1733  
 1734  
 1735  
 1736  
 1737  
 1738  
 1739  
 1740  
 1741  
 1742  
 1743  
 1744  
 1745  
 1746  
 1747  
 1748  
 1749  
 1750  
 1751  
 1752  
 1753  
 1754  
 1755  
 1756  
 1757  
 1758  
 1759  
 1760  
 1761  
 1762  
 1763  
 1764  
 1765  
 1766  
 1767  
 1768  
 1769  
 1770  
 1771  
 1772  
 1773  
 1774  
 1775  
 1776  
 1777  
 1778  
 1779  
 1780  
 1781  
 1782  
 1783  
 1784  
 1785  
 1786  
 1787  
 1788  
 1789  
 1790  
 1791  
 1792  
 1793  
 1794  
 1795  
 1796  
 1797  
 1798  
 1799  
 1800  
 1801  
 1802  
 1803  
 1804  
 1805  
 1806  
 1807  
 1808  
 1809  
 1810  
 1811  
 1812  
 1813  
 1814  
 1815  
 1816  
 1817  
 1818  
 1819  
 1820  
 1821  
 1822  
 1823  
 1824  
 1825  
 1826  
 1827  
 1828  
 1829  
 1830  
 1831  
 1832  
 1833  
 1834  
 1835  
 1836  
 1837  
 1838  
 1839  
 1840  
 1841  
 1842  
 1843  
 1844  
 1845  
 1846  
 1847  
 1848  
 1849  
 1850  
 1851  
 1852  
 1853  
 1854  
 1855  
 1856  
 1857  
 1858  
 1859  
 1860  
 1861  
 1862  
 1863  
 1864  
 1865  
 1866  
 1867  
 1868  
 1869  
 1870  
 1871  
 1872  
 1873  
 1874  
 1875  
 1876  
 1877  
 1878  
 1879  
 1880  
 1881  
 1882  
 1883  
 1884  
 1885  
 1886  
 1887  
 1888  
 1889  
 1890  
 1891  
 1892  
 1893  
 1894  
 1895  
 1896  
 1897  
 1898  
 1899  
 1900  
 1901  
 1902  
 1903  
 1904  
 1905  
 1906  
 1907  
 1908  
 1909  
 1910  
 1911  
 1912  
 1913  
 1914  
 1915  
 1916  
 1917  
 1918  
 1919  
 1920  
 1921  
 1922  
 1923  
 1924  
 1925  
 1926  
 1927  
 1928  
 1929  
 1930  
 1931  
 1932  
 1933  
 1934  
 1935  
 1936  
 1937  
 1938  
 1939  
 1940  
 1941  
 1942  
 1943  
 1944  
 1945  
 1946  
 1947  
 1948  
 1949  
 1950  
 1951  
 1952  
 1953  
 1954  
 1955  
 1956  
 1957  
 1958  
 1959  
 1960  
 1961  
 1962  
 1963  
 1964  
 1965  
 1966  
 1967  
 1968  
 1969  
 1970  
 1971  
 1972  
 1973  
 1974  
 1975  
 1976  
 1977  
 1978  
 1979  
 1980  
 1981  
 1982  
 1983  
 1984  
 1985  
 1986  
 1987  
 1988  
 1989  
 1990  
 1991  
 1992  
 1993  
 1994  
 1995  
 1996  
 1997  
 1998  
 1999  
 2000  
 2001  
 2002  
 2003  
 2004  
 2005  
 2006  
 2007  
 2008  
 2009  
 2010  
 2011  
 2012  
 2013  
 2014  
 2015  
 2016  
 2017  
 2018  
 2019  
 2020  
 2021  
 2022  
 2023  
 2024  
 2025  
 2026  
 2027  
 2028  
 2029  
 2030  
 2031  
 2032  
 2033  
 2034  
 2035  
 2036  
 2037  
 2038  
 2039  
 2040  
 2041  
 2042  
 2043  
 2044  
 2045  
 2046  
 2047  
 2048  
 2049  
 2050  
 2051  
 2052  
 2053  
 2054  
 2055  
 2056  
 2057  
 2058  
 2059  
 2060  
 2061  
 2062  
 2063  
 2064  
 2065  
 2066  
 2067  
 2068  
 2069  
 2070  
 2071  
 2072  
 2073  
 2074  
 2075  
 2076  
 2077  
 2078  
 2079  
 2080  
 2081  
 2082  
 2083  
 2084  
 2085  
 2086  
 2087  
 2088  
 2089  
 2090  
 2091  
 2092  
 2093  
 2094  
 2095  
 2096  
 2097  
 2098  
 2099  
 2100  
 2101  
 2102  
 2103  
 2104  
 2105  
 2106  
 2107  
 2108  
 2109  
 2110  
 2111  
 2112  
 2113  
 2114  
 2115  
 2116  
 2117  
 2118  
 2119  
 2120  
 2121  
 2122  
 2123  
 2124  
 2125  
 2126  
 2127  
 2128  
 2129  
 2130  
 2131  
 2132  
 2133  
 2134  
 2135  
 2136  
 2137  
 2138  
 2139  
 2140  
 2141  
 2142  
 2143  
 2144  
 2145  
 2146  
 2147  
 2148  
 2149  
 2150  
 2151  
 2152  
 2153  
 2154  
 2155  
 2156  
 2157  
 2158  
 2159  
 2160  
 2161  
 2162  
 2163  
 2164  
 2165  
 2166  
 2167  
 2168  
 2169  
 2170  
 2171  
 2172  
 2173  
 2174  
 2175  
 2176  
 2177  
 2178  
 2179  
 2180  
 2181  
 2182  
 2183  
 2184  
 2185  
 2186  
 2187  
 2188  
 2189  
 2190  
 2191  
 2192  
 2193  
 2194  
 2195  
 2196  
 2197  
 2198  
 2199  
 2200  
 2201  
 2202  
 2203  
 2204  
 2205  
 2206  
 2207  
 2208  
 2209  
 2210  
 2211  
 2212  
 2213  
 2214  
 2215  
 2216  
 2217  
 2218  
 2219  
 2220  
 2221  
 2222  
 2223  
 2224  
 2225  
 2226  
 2227  
 2228  
 2229  
 2230  
 2231  
 2232  
 2233  
 2234  
 2235  
 2236  
 2237  
 2238  
 2239  
 2240  
 2241  
 2242  
 2243  
 2244  
 2245  
 2246  
 2247  
 2248  
 2249  
 2250  
 2251  
 2252  
 2253  
 2254  
 2255  
 2256  
 2257  
 2258  
 2259  
 2260  
 2261  
 2262  
 2263  
 2264  
 2265  
 2266  
 2267  
 2268  
 2269  
 2270  
 2271  
 2272  
 2273  
 2274  
 2275  
 2276  
 2277  
 2278  
 2279  
 2280  
 2281  
 2282  
 2283  
 2284  
 2285  
 2286  
 2287  
 2288  
 2289  
 2290  
 2291  
 2292  
 2293  
 2294  
 2295  
 2296  
 2297  
 2298  
 2299  
 2300  
 2301  
 2302  
 2303  
 2304  
 2305  
 2306  
 2307  
 2308  
 2309  
 2310  
 2311  
 2312  
 2313  
 2314  
 2315  
 2316  
 2317  
 2318  
 2319  
 2320  
 2321  
 2322  
 2323  
 2324  
 2325  
 2326  
 2327  
 2328  
 2329  
 2330  
 2331  
 2332  
 2333  
 2334  
 2335  
 2336  
 2337  
 2338  
 2339  
 2340  
 2341  
 2342  
 2343  
 2344  
 2345  
 2346  
 2347  
 2348  
 2349  
 2350  
 2351  
 2352  
 2353  
 2354  
 2355  
 2356  
 2357  
 2358  
 2359  
 2360  
 2361  
 2362  
 2363  
 2364  
 2365  
 2366  
 2367  
 2368  
 2369  
 2370  
 2371  
 2372  
 2373  
 2374  
 2375  
 2376  
 2377  
 2378  
 2379  
 2380  
 2381  
 2382  
 2383  
 2384  
 2385  
 2386  
 2387  
 2388  
 2389  
 2390  
 2391  
 2392  
 2393  
 2394  
 2395  
 2396  
 2397  
 2398  
 2399  
 2400  
 2401  
 2402  
 2403  
 2404  
 2405  
 2406  
 2407  
 2408  
 2409  
 2410  
 2411  
 2412  
 2413  
 2414  
 2415  
 2416  
 2417  
 2418  
 2419  
 2420  
 2421  
 2422  
 2423  
 2424  
 2425  
 2426  
 2427  
 2428  
 2429  
 2430  
 2431  
 2432  
 2433  
 2434  
 2435  
 2436  
 2437  
 2438  
 2439  
 2440  
 2441  
 2442  
 2443  
 2444  
 2445  
 2446  
 2447  
 2448  
 2449  
 2450  
 2451  
 2452  
 2453  
 2454  
 2455  
 2